

PLEASE NOTE

**This image contains more than one label
approved for this product on this date.**

ACCEPTED

NET CONTENTS

POUNDS

T-H

CASORON W-50

MAR 20 1964
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 148-615

ACTIVE INGREDIENT:

Dichlobenil*.....	50.0%
INERT INGREDIENTS.....	50.0%
	100.0%

*2,6-dichlorobenzonitrile

A SELECTIVE PRE-EMERGENCE HERBICIDE FOR CONTROL OF CRABGRASS, PIGWEED AND CERTAIN OTHER ANNUAL WEEDS.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

T-H CASORON W-50 is recommended for control of certain grassy and broadleaved weeds such as crabgrass, redroot pigweed, henbit, smartweed, wild mustard, pineapple weed, knotweed, chickweed, shepherds purse, lambsquarters, groundsel and foxtail. Apply 8 to 12 pounds of T-H CASORON W-50 per acre, based on area actually sprayed, in sufficient water for uniform coverage (at least 50 gallons per acre). In Florida, apply 8 to 16 pounds per acre. If treatment is followed by application of 1/2 to 1 inch of water by overhead irrigation, the lower rates are recommended. Apply T-H CASORON W-50 only to prepared, weed-free soil either in early spring before seeds of annual weeds germinate or after cultivation has removed growing weeds.

NURSERY STOCK: T-H CASORON W-50 may be used for weed control on the plants listed below. CASORON W-50 is for field use only. Do not apply within 4 weeks of transplanting.

ORNAMENTALS

Arbor Vitae
 Boxwood (Buxus spp.)
 Holly (Ilex spp.
 except Crenata)

Juniper
 (Juniperus spp.)
 Pittosporum

Roses
 Andromeda (Pieris
spp.)

Yew (Taxus spp.)
 Rhododendron (not Azaleas)
 Pine (Pinus spp.)
 Elwood Cypress
 (Chamaecyparis spp.)

Podocarpus
 Pyracantha
 Forsythia
 Privet (Ligustrum spp.)
 Euonymus
 Barberry

Ivy (Hedera halix)
 Fir (Abies spp.)
 Spruce (Picea spp.)
 Cotoneaster
 Peony
 Gladiolus

PRECAUTIONS

Continuous agitation within the spray tank is essential. Wash out spray tank thoroughly after use. Do not use in seed beds, transplant or cutting beds. Store in a tightly closed container in a dry place.

CAUTION

HARMFUL IF SWALLOWED! Do not breathe dust. Do not allow contact with eyes or skin. Use with caution. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

NOTICE

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely neither Thompson-Hayward Chemical Company nor seller makes any warranty or representation expressed or implied concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. Neither Thompson-Hayward Chemical Company nor seller shall be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this

label. Damages, if any, which may result from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages such as those resulting from crop injury, or failure to give adequate control.

PHILIPS ROXANE DIV.

THOMPSON-HAYWARD CHEMICAL COMPANY

SUBSIDIARY OF PHILIPS ELECTRONICS AND PHARMACEUTICAL INDUSTRIES CORP.

KANSAS CITY, MISSOURI 64141

NEXT

LABEL

NET CONTENTS

POUNDS

T-H

CASORON W-50

ACCEPTED
MAR 20 1964
<small>UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTER- ED UNDER NO. 148-615</small>

ACTIVE INGREDIENT:

Dichlobenil*.....	50.0%
INERT INGREDIENTS.....	50.0%
	100.0%

*2,6-dichlorobenzonitrile

A SELECTIVE PRE-EMERGENCE HERBICIDE FOR CONTROL CRABGRASS, PIGWEED AND CERTAIN OTHER ANNUAL WEEDS.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

T-H CASORON W-50 is recommended for control of certain grassy and broadleaved weeds such as crabgrass, redroot pigweed, henbit, smartweed, wild mustard, pineapple weed, knotweed, chickweed, shepherds purse, lambsquarters, groundsel and foxtail. Apply 8 to 12 pounds of T-H CASORON W-50 per acre, based on area actually sprayed, in sufficient water for uniform coverage (at least 50 gallons per acre). In Florida, apply 8 to 16 pounds per acre. If treatment is followed by application of 1/2 to 1 inch of water by overhead irrigation, the lower rates are recommended. Apply T-H CASORON W-50 only to prepared, weed-free soil either in early spring before seeds of annual weeds germinate or after cultivation has removed growing weeds.

FRUIT (Non-bearing) AND FRUIT NURSERIES

T-H CASORON W-50 may be used for weed control on the plants listed below. CASORON W-50 is for field use only. Do not apply within 4 weeks of transplanting. On orchard trees of bearing age, do not make application during fall or spring preceding fruit harvest.

- | | |
|--------|--------|
| Apple | Plum |
| Peach | Cherry |
| Citrus | |

PRECAUTIONS

Continuous agitation within the spray tank is essential. Wash out spray tank thoroughly after use. Do not use in seed beds, transplant or cutting beds. Store in a tightly closed container in a dry place.

CAUTION

HARMFUL IF SWALLOWED! Do not breathe dust. Do not allow contact with eyes or skin. Use with caution. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

NOTICE

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely neither Thompson-Hayward Chemical Company nor seller makes any warranty or representation expressed or implied concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. Neither Thompson-Hayward Chemical Company nor seller shall be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this label. Damages, if any, which may result from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages such as those resulting from crop injury, or failure to give adequate control.

PHILIPS ROXANE DIV.

THOMPSON-HAYWARD CHEMICAL COMPANY

SUBSIDIARY OF PHILIPS ELECTRONICS AND PHARMACEUTICAL INDUSTRIES CORP.

KANSAS CITY, MISSOURI 64141

NEXT

LABEL


NET CONTENTS

POUNDS

T-H

CASORON W-50

ACTIVE INGREDIENT:	
Dichlobenil*.....	50.0%
INERT INGREDIENTS.....	50.0%
	100.0%



*2,6-dichlorobenzonitrile

A SELECTIVE PRE-EMERGENCE HERBICIDE FOR CONTROL OF CRABGRASS, PIGWEED AND CERTAIN OTHER ANNUAL WEEDS.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

T-H CASORON W-50 is recommended for control of certain grassy and broad-leaved weeds such as crabgrass, redroot pigweed, henbit, smartweed, wild mustard, pineapple weed, knotweed, chickweed, shepherds purse, lambsquarters, groundsel and foxtail. Apply 8 to 12 pounds of T-H CASORON W-50 per acre, based on area actually sprayed, in sufficient water for uniform coverage (at least 50 gallons per acre). Apply T-H CASORON W-50 only to prepared, weed-free soil either in early spring before seeds of annual weeds germinate or after cultivation has removed growing weeds.

SHELTERBELTS AND FOREST PLANTINGS: T-H CASORON W-50 may be used for weed control on the plants listed below. CASORON W-50 is for field use only. Do not apply within 4 weeks of transplanting. May be used for maintenance weed control.

Jack Pine
Cottonwood
American Elm
Chinese Elm
Hackberry
Caragana

Boxelder
Russian Olive
Honey Locust
Green Ash
Eastern Red Cedar

PRECAUTIONS

Continuous agitation within the spray tank is essential. Wash out spray tank thoroughly after use. Do not use in seed beds, transplant or cutting beds. Store in a tightly closed container in a dry place.

CAUTION

HARMFUL IF SWALLOWED! Do not breathe dust. Do not allow contact with eyes or skin. Use with caution. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

NOTICE

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely neither Thompson-Hayward Chemical Company nor seller makes any warranty or representation expressed or implied concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. Neither Thompson-Hayward Chemical Company nor seller shall be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this label. Damages, if any, which may result from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages such as those resulting from crop injury, or failure to give adequate control.

PHILIPS ROXANE DIV.

THOMPSON-HAYWARD CHEMICAL COMPANY

SUBSIDIARY OF PHILIPS ELECTRONICS AND PHARMACEUTICAL INDUSTRIES CORP.

KANSAS CITY, MISSOURI 64141

NEXT

LABEL

148-615

03/20/1964

ACCEPTED

MAR 20 1964

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 148-615

POUNDS

1/3

NET CONTENTS

T-H

CASORON W-50

ACTIVE INGREDIENT:

Dichlobenil*.....	50.0%
INERT INGREDIENTS.....	50.0%
	100.0%

*2,6-dichlorobenzonitrile

A SELECTIVE PRE-EMERGENCE HERBICIDE FOR CONTROL OF CRABGRASS, PIGWEED AND CERTAIN OTHER ANNUAL WEEDS.

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY.

T-H CASORON W-50 is recommended for control of certain grassy and broadleaved weeds such as crabgrass, redroot pigweed, wild mustard, knotweed, chickweed, shepherds purse, and lambsquarters. Use sufficient water for uniform coverage, at least 50 gallons per acre.

BERMUDAGRASS ESTABLISHMENT: CASORON W-50 spray should be applied immediately after setting Bermudagrass sprigs in well-prepared, weed-free soil. Apply 4 to 8 pounds of T-H CASORON W-50 per acre, based on area actually sprayed. Application may be made as a band treatment over newly-set sprigs if planted in rows or as a broadcast overall treatment. Do not allow animals to graze treated areas, do not use crop as animal feed. In Southern New Jersey and adjoining Eastern Coastal areas use the lower rate of application. In other Bermudagrass areas the higher rate may result in more efficient weed control.

WEED CONTROL IN ESTABLISHED BERMUDAGRASS TURF: To control annual weeds in Bermudagrass turf on airports, cemeteries and recreational (non-crop) areas such as football fields, baseball fields and golf courses, apply 8 to 12

pounds of T-H CASORON W-50 per acre prior to weed seed germination. Do not allow animals to graze treated areas, do not use crop as animal feed.

PRECAUTIONS

Continuous agitation within the spray tank is essential. Wash out spray tank thoroughly after use. Do not apply more than once per year. Store in a tightly closed container in a dry place.

CAUTION

HARMFUL IF SWALLOWED! Do not breathe dust. Do not allow contact with eyes or skin. Use with caution. Avoid contamination of feed and foodstuffs. Keep out of reach of children.

NOTICE

The above directions are based upon use under normal and reasonably foreseeable conditions. If the directions are followed and precautions on this label observed, available research indicates that this product will give satisfactory results. However, because climatic, geographic, and other conditions of use will vary widely neither Thompson-Hayward Chemical Company nor seller makes any warranty or representation expressed or implied concerning this product or its use except that it conforms to the chemical description on the label and no person is authorized to make any other representations upon their behalf. Neither Thompson-Hayward Chemical Company nor seller shall be held responsible for damages resulting from the handling, storage, or use of this product for any reason except the failure of the material to conform to the chemical description on this label. Damages, if any, which may result from the failure of this product to conform to the description on this label shall be limited to direct damages but shall not include consequential damages such as those resulting from crop injury, or

failure to give adequate control.

PHILIPS ROXANE DIV.

THOMPSON-HAYWARD CHEMICAL COMPANY

SUBSIDIARY OF PHILIPS ELECTRONICS AND PHARMACEUTICAL INDUSTRIES CORP.

KANSAS CITY, MISSOURI 64141
