

**MIXING INSTRUCTIONS**

Determine the amount of Ded-Weed LV-6D to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed LV-6D recommended per acre. Add this amount of Ded-Weed LV-6D to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

For control of resistant species of brush and certain timberland applications, use other Thompson-Hayward products such as Ded-Weed LV-2D2T and Ded-Weed LV-4T.


**FOR DILUTION CALCULATIONS—**

1 pint contains . . . lb 2,4-D Acid

**PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED LV-6D**

- Carpet Weed      Mustards
- Cocklebur        Proweed
- Lamb's Quarters   Wild Morning Glory

If stored below freezing temperatures it may be necessary to warm to 40° F. and agitate thoroughly before using.



**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



**2,4-D 6 POUNDS PER GALLON  
LOW VOLATILE ESTER**

**KILLS WEEDS IN PASTURES, CROPS AND NON-CROP  
AREAS, FENCE ROWS AND RIGHT-OF-WAYS**

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN  
SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.**

**ACTIVE INGREDIENT:**

2,4-dichlorophenoxyacetic acid isooctyl ester . . . . . 93.5%

**INERT INGREDIENTS:**

Total . . . . . 100.0%

\*(Equivalent to 2,4-dichlorophenoxyacetic acid—62%)

EPA Reg No 148-616

CG-73-A

EPA Est No 148-KS-1

**NET CONTENTS      GAL. (            litres)**



**P.O. BOX 2383, KANSAS CITY, KANSAS 66110**

### CAUTION

Avoid contact with skin, eyes or clothing. Harmful if swallowed.

Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purposes.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Although this product has been formulated with low volatile esters in order to reduce the hazard of vapor damage. Vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, moisture and wind movements are ideal for vaporization and drift of the vapor.

Do not re-use this container. Destroy by burying in a safe place.

Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides or other material to susceptible crops.

Do not store near fertilizers, seed, insecticides, fungicides or food stuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover lawns.

Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.

### DIRECTIONS BEFORE USE READ DIRECTIONS STATEMENTS OF WEED CONTROL

WEEDS	Amount Ded Weed LV-6D Per Acre	Amount 2,4-D Acid Per Acre	
BROADLEAF WEEDS IN			
SPRING WHEAT, RYE, BARLEY	1.5 to 1.2 pint	1.4 to 1.3 lb	Apply full after milk
OATS	1.3 pint	1.4 lb	
FALL WHEAT	1.3 to 2.3 pint	1.4 to 1.2 lb	
FIELD CORN PRE-EMERGENCE	1.1 to 2 pints	1 to 1.2 lbs	Any time after 2 to 5 days
CORN, SORGHUMS POST-EMERGENCE	1.4 to 1.3 pint	3.1 to 1.4 lb	When weed crop is 4 to 6 inches high
PASTURES, RANGELAND	1 pint	3.4 lb	When annual weeds are green perennials in seed

\*Do not graze dairy animals on treated areas within 7 days after heading begins. Do not apply when grass is in boot to milk.

### FOR PERENNIAL WEEDS\* IN PASTURES AND NON-CROP LANDS SUCH AS DRAINAGE DITCH BANKS

WEEDS	Amount Ded Weed LV-6D Per Acre	Amount 2,4-D Acid Per Acre	
CANADA THISTLE, AND PERENNIAL SOW THISTLE	1 pint	3.4 lb	Apply during again when
BINDWEED	1 pint	3.4 lb	Apply bud to
WHITE TOP (HOARY CRESS)	1 pint	3.4 lb	In spring when and foliage
RUSSIAN KNAPWEED	1 pint	4 lb	In spring be

\*Hard-to-kill weeds such as Canadian Thistle may not be controlled. Usually will be prevented. Repeated treatments generally give control.

148-6100

5%  
5%  
0%  
KS-1  
S)  
10

## CAUTION

Avoid contact with skin, eyes, or clothing. Harmful if swallowed.

Avoid contamination of water supplies that may be used to irrigate or water susceptible crops, or used for domestic purpose.

Under no circumstances should this herbicide product or any 2,4-D or 2,4,5-T weed killer be used in the vicinity of cotton, tomatoes, garden crops, grapes, ornamentals, or other susceptible crops, as severe damage may result. Do not apply on a windy day. Extreme care must be exercised to prevent drifting during application. (Coarse sprays are less likely to drift.)

Although this product has been formulated with low volatile esters in order to reduce the hazard of vapor damage, vapors resulting from the volatilization (evaporation) of this product after application may cause injury to susceptible plants in the immediate vicinity or for great distances if temperature, moisture and wind movements are ideal for vaporization and drift of the vapor.

Do not re-use this container. Destroy by burying in a safe place.

Do not use equipment used in applying this product or any 2,4-D or 2,4,5-T weed killer to apply insecticides, fungicides, or other material to susceptible crops.

Do not store near fertilizers, seed, insecticides, fungicides or food stuffs. May cause injury to bentgrass, St. Augustine grass, Dichondra, carpetgrass or clover lawns.

Do not use or store near heat or open flame. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops.

## DIRECTIONS BEFORE USE READ DIRECTIONS, CAUTION AND STATEMENTS CAREFULLY WEED CONTROL IN CROPS

WEEDS IN	Amount Ded Weed LV 50 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION
BROADLEAF WEEDS IN			
SPRING WHEAT, RYE, BARLEY	1 pint	1.4 lb	Apply for control of broadleaf weeds after milk stage to the next
OLDS	1 pint	1.4 lb	
FALL WHEAT	1 pint	1.4 lb	
FIELD CORN PRE EMERGENCE	1 pint	1.4 lb	Any time after planting, preferably 3 to 5 days, but before emergence
CORN, SORGHUMS POST EMERGENCE	1 pint	1.4 lb	When weeds are 1 to 2 inches tall, crops 4 to 18 inches tall
PASTURES RANGELAND		1.4 lb	When annual weeds are 1 to 2 inches tall, perennial weeds are 1 to 2 inches tall, or near bud stage

\*Do not graze dairy animals on treated areas within 7 days after application. Do not apply when grass is in boot to milk stage.

## FOR PERENNIAL WEEDS\* IN PASTURE, RANGELAND AND NON-CROP LANDS SUCH AS FENCE ROWS, DRAINAGE DITCH BANKS, RIGHT-OF-WAYS

WEEDS	Amount Ded Weed LV 50 Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION
CANADA THISTLE AND PERENNIAL SOW THISTLE	1 pint	1.4 lb	Apply during early bud stage and repeat when plants are formed
BINDWEED	1 pint	1.4 lb	Apply during bud stage
WHITE TOP (HOARY CRESS)	1 pint	1.4 lb	Apply when plants are 1 to 2 inches tall and buds are enlarged
RUSSIAN KNAPWEED		1.4 lb	

\*Hard to kill weeds such as Canadian Thistle may not be controlled by one application. Control will be prevented. Repeated treatments generally give good control.

148-616