

NET CONTENTS

ACCEPTED
148-677



CYTHION LV CONCENTRATE

GALLON

ACCEPTED
24, 1977
148-677

For use only in spray tanks...
148-677 PUMP

BY-PASS

When pump is repaired...
CYTHION LV CONCENTRATE

BOOM AND NOZZLE PLACEMENT

The distance between the left and right...
nozzles should be 4 to 6 feet

NOZZLES

Use only 4 to 6 foot diameter nozzles...
with a spray angle of 40 to 60 degrees

When using a spray pump...
the nozzle should be 4 to 6 feet from the crop

When using a backpack sprayer...
the nozzle should be 4 to 6 feet from the crop

CYTHION LV CONCENTRATE

Contains 10% active ingredient...
100% active ingredient

DISPOSAL OF DRUMS

Drums should be cleaned...
and disposed of properly

CAUTION! Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS

**HARMFUL BY SWALLOWING,
INHALATION OR SKIN CONTACT**

Avoid Breathing Spray Mist

Avoid Contact With Skin

Wash Thoroughly After Handling

Change Contaminated Clothing

Do Not Store Near Feed or Foodstuffs

Cythion may be toxic to certain species of fish.

Avoid contamination of lakes, ponds and streams.

ACTIVE INGREDIENT

Methidathion (O,O Dimethyl dithio-phosphate)
of diethylene glycol dimethyl ether 95

INERT INGREDIENTS

5
100

*O,O dimethyl phosphorothioate of diethyl mercaptosuccinate
and diethylene glycol dimethyl ether

CYTHION - Trademark of American Cyanamid Company

DIRECTIONS FOR USE

CYTHION LV CONCENTRATE is a liquid undiluted for a truck
or other aerial application...
at altitudes of 20 to 25 feet

NOTE: CYTHION LV CONCENTRATE may cause spitting on
skin. Do not permit spray to come in contact with face...
If an accidental exposure does occur, the person should wash immediately.

IMPORTANT

When using a spray pump...
the nozzle should be 4 to 6 feet from the crop

1. Adjust pump pressure to 100 psi or more.
2. Drive slowly over the field with an average speed of 2 to 3 miles per hour.
3. Keep nozzle at a constant height of 4 to 6 feet above the crop.
4. Maintain a constant speed of 4 to 6 miles per hour.
5. Maintain a constant spray rate of 10 to 15 gallons per acre.
6. Do not use in windy conditions.
7. Do not use in cold weather.
8. Do not use on wet crops.
9. Do not use on crops that are stressed.
10. Do not use on crops that are dormant.

THOMPSON-HAYWARD CHEMICAL COMPANY
P. O. Box 2793 • Kansas City, Kansas 66110

CROP	PEST	FLUID OUNCE PER ACRE	INTERVAL BETWEEN APPLICATION AND HARVEST
ALFALFA, CLOVER PASTURE AND RANGE, GRASS, GRASS, GRASS, HAY, NONAGRICULTURAL LAND (WASTE LANDS, ROADSIDES, SOIL BANK LANDS)			
WHEAT			
OATS			
WHEAT			
OATS			
COTTON			
PEANUTS			
SUGAR BEETS			
CORN			
BEANS			
NONAGRICULTURAL LANDS			
FREE CATTLE FEED LOTS AND BARRIERS			

NOTICE: This product is...
Do not use on...
Do not use on...