

DE-PESTER®

THIO 3% DUST

RECOMMENDED USES

ARTICHOKES: USE 30 TO 50 LBS. PER ACRE TO CONTROL ARTICHOKE APHID AND GREEN PEACH APHID. REPEAT AS NEEDED FOR CONTROL. DO NOT APPLY AFTER BUDS START TO FORM.

BEANS: USE 20 TO 30 LBS. PER ACRE TO CONTROL MEXICAN BEAN BEETLE. DO NOT APPLY AFTER PODS BEGIN TO FORM. DO NOT FEED TREATED THRESHINGS TO LIVESTOCK.

BROCCOLI, CABBAGE & CAULIFLOWER: USE 25 LBS. PER ACRE TO CONTROL CABBAGE LOOPER, IMPORTED CABBAGE WORM, DIAMONDBACK MOTM LARVAE, CROSS-STRIPED CABBAGE WORM, FLEA BEETLE, HARLEQUIN BUG AND STINK BUGS. REPEAT AS NECESSARY TO MAINTAIN CONTROL. DO NOT APPLY TO CAULIFLOWER AFTER EDIBLE PARTS START TO FORM. DO NOT APPLY TO CAULIFLOWER AFTER EDIBLE PARTS START TO FORM. DO NOT APPLY TO BROCCOLI AND CABBAGE WITHIN 7 DAYS OF HARVEST.

COTTON: USE 30 TO 50 LBS. PER ACRE TO CONTROL STINK BUGS, LYGUS BUGS, BOLL-WORM AND CABBAGE LOOPER. REPEAT AS NECESSARY TO MAINTAIN CONTROL. DO NOT APPLY AFTER BOLLS BEGIN TO OPEN. DO NOT GRAZE MEAT OR DAIRY ANIMALS IN TREATED FIELDS.

CUCUMBERS, MELONS, PUMPKINS, SUMMER AND WINTER SQUASH: USE 25 TO 30 LBS. PER ACRE TO CONTROL APHIDS. REPEAT AS NECESSARY TO MAINTAIN CONTROL. MAY BE APPLIED UP TO DAY OF HARVEST.

EGGPLANTS: USE 20 LBS. PER ACRE TO CONTROL APHIDS, FLEA BEETLES, COLORADO POTATO BEETLE, AND WHITEFLY. REPEAT AS NECESSARY TO MAINTAIN CONTROL. DO NOT APPLY WITHIN 7 DAYS OF HARVEST.

GRAPES: USE 30 TO 50 LBS. PER ACRE TO CONTROL GRAPE LEAFHOPPER. DO NOT APPLY WITHIN 7 DAYS OF HARVEST. DO NOT USE ON CONCORD VARIETY AS INJURY IS LIKELY TO OCCUR.

PEAS (SEED) USE 35 TO 40 LBS. PER ACRE TO CONTROL PEA APHID AND PEA WEEVIL. DO NOT FEED TREATED VINES OR THRESHINGS TO DAIRY OR MEAT ANIMALS.

POTATOES: USE 30 TO 50 LBS. PER ACRE TO CONTROL FLEA BEETLE, COLORADO BEETLE, LEAFHOPPERS, APHIDS, POTATO TUBERWORM, LEAF-FOOTED PLANT BUG, SOUTHERN ARMY WORM, GREEN STINK BUG, POTATO PSYLLID AND EUROPEAN CORN BORER. APPLY AS NEED TO HARVEST.

ACTIVE INGREDIENT:

ENDOSULFAN*.....3.0%
INERT INGREDIENTS.....97.0%

TOTAL.....100.0%

*HEXACHLOROHYDROMETHANO-2, 4, 3-BENZO-DIOXATHIEPIN OXIDE

RECOMMENDED USES (CONTINUED)

PEPPERS & TOMATOES: USE 25 TO 30 LBS. PER ACRE TO CONTROL COLORADO POTATO BEETLE, FLEA BEETLE, APHIDS, AND WHITEFLY. REPEAT AS NECESSARY TO MAINTAIN CONTROL. DO NOT APPLY WITHIN 1 DAY OF PICKING.

WALNUTS: USE 50 TO 60 LBS. PER ACRE TO CONTROL WALNUT APHIDS. REPEAT AS REQUIRED TO MAINTAIN CONTROL. DO NOT APPLY AFTER HUSK SPLIT.

THIS PRODUCT WILL KILL FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. CONTAINERS SHOULD BE DISPOSED BY BURNING (KEEP OUT OF SMOKE).

CAUTION

KEEP OUT OF REACH OF CHILDREN !

SUGGESTIONS FOR USE

APPLY WHEN INSECTS ARE ON PLANTS AND WHEN AIR IS QUIET, PREFERABLY IN EARLY MORNING. DO NOT MIX OR DILUTE THIS DUST WITH OTHER MATERIALS. EVEN COVERAGE OF INFESTED PLANTS IS ESSENTIAL FOR CONTROL. DOSAGE PER ACRE WILL DEPEND ON SIZE OF INFESTATION. USE ONLY SUFFICIENT MATERIAL TO EFFECT DESIRABLE CONTROL.

CAUTION

HARMFUL IF SWALLOWED, INHALED, OR ABSORBED THROUGH THE SKIN, EYES, AND CLOTHING. AVOID INHALATION OR CONTACT WITH SKIN, EYES, AND CLOTHING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. IN CASE OF CONTACT WITH SKIN OR EYES, FLUSH WITH PLENTY OF WATER; FOR EYES GET MEDICAL ATTENTION. WASH WITH SOAP AND WATER AFTER HANDLING AND BEFORE EATING OR SMOKING; WEAR CLEAN CLOTHING. DO NOT APPLY OR ALLOW TO DRIFT TO AREAS OCCUPIED BY UNPROTECTED HUMANS OR BENEFICIAL ANIMALS.

WORKERS ENTERING TREATED AREAS THE DAY OF APPLICATION SHOULD WEAR PROTECTIVE CLOTHING.

DO NOT ALLOW THIS PRODUCT TO DRIFT FROM AREA BEING TREATED.

ACCEPTED

JUL 20 1967

UNDER THE FEDERAL INSECTICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER NO. 148-698 TO ATTACHED COMMENTS

THOMPSON-HAYWARD CHEMICAL CO.

FRESNO, CALIFORNIA — KANSAS CITY, MISSOURI

NOTICE: No warranty, expressed or implied, is made concerning this product or its use that it conforms to the chemical description on this label and no person is authorized to make any other representation. Neither Thompson-Hayward Chemical Company nor seller shall be liable for any damages resulting from handling, storage or use of this product.

USDA Reg. No. 148-698