



PHOSDRIN* 2% DUST

DANGER

POISON

WARNING

DIRECTIONS FOR USE

BEFORE APPLICATION WHEN INSECTS FIRST APPEAR AND REPEAT AS OFTEN AS NECESSARY TO MAINTAIN CONTROL. APPLY SO AS TO GIVE UNIFORM COVERAGE.

RECOMMENDED USES

ALFALFA, CLOVER, OIL AND SORGHUM: FOR CONTROL OF APHIDS, SPIDER MITES, LEAFHOPPERS, GRASSHOPPERS, LYGUS BUGS, ALFALFA CATERPILLAR, ALFALFA WEEVIL LARVAE, CORN EARWORM (ON SORGHUM) AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY ON ALFALFA, CLOVER AND CORN WITHIN 1 DAY OF HARVEST, AND ON SORGHUM WITHIN 3 DAYS OF HARVEST.

ARTICHOKES: FOR CONTROL OF APHIDS AND ARTICHOKE FLY. BOTH APPLY 15-50 LBS. PER ACRE. FOR FLY MOths USE HINER LOGANES. DO NOT APPLY WITHIN 2 DAYS OF HARVEST.

BEANS AND PEAS: FOR CONTROL OF APHIDS, SPIDER MITES, LEAFHOPPERS, MEXICAN BEAN BEETLE, GRASSHOPPERS AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY WITHIN 1 DAY OF HARVEST.

BELTS, LETTUCE, ONIONS AND SPINACH: FOR CONTROL OF APHIDS, SPIDER MITES, DIPTEROUS LEAF-MINER (ADULTS), THRIPS, LYGUS BUGS, FALSE CHINCH BUG, LEAFHOPPERS, GRASSHOPPERS, CORN EARWORM, CABBAGE LOOPER, IMPORTED CABBAGEWORM, SALT-MARSH CATERPILLAR AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY ON LETTUCE WITHIN 2 DAYS OF HARVEST; ON CARA AND ONIONS WITHIN 1 DAY OF HARVEST; ON SPINACH WITHIN 1 DAY OF HARVEST; AND ON BELTS WITHIN 3 DAYS OF HARVEST.

CELERY AND CUCUMBERS: FOR CONTROL OF APHIDS, SPIDER MITES, DIPTEROUS LEAF-MINER (ADULTS), LYGUS BUGS, LEAFHOPPERS, GRASSHOPPERS, CABBAGE LOOPER, SALT-MARSH CATERPILLAR AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY ON CABBAGES WITHIN 1 DAY OF HARVEST AND ON CELERY WITHIN 3 DAYS OF HARVEST.

CUCUMBERS, BRUSSELS SPROUTS, CABBAGE, CABBAGE FLOWER, CUCUMBERS, CABBAGE, STARD GREENS AND TURNIPS: FOR CONTROL OF APHIDS, SPIDER MITES, DIPTEROUS LEAF-MINER (ADULTS), LYGUS BUGS, LEAFHOPPERS, GRASSHOPPERS, CABBAGE LOOPER, IMPORTED CABBAGEWORM, SALT-MARSH CATERPILLAR AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY ON BRUSSELS AND CABBAGE WITHIN 1 DAY OF HARVEST AND ON BRUSSELS SPROUTS, CABBAGE FLOWER,

ACTIVE INGREDIENTS:

ALPHA ISOMER OF 2-CARBOETHYL-1-METHYLVINYL DIMETHYL PHOSPHATE ..	1.2%..
RELATED COMPOUNDS	0.6%..
INERT INGREDIENTS	98.2%
TOTAL	100.0%

- REGISTERED T. M. SHELL CHEMICAL COMPANY
- EQUIVALENT TO 2.0% PHOSDRIN INSECTICIDE

RECOMMENDED USES (CONTINUED)

CUCUMBERS (CONTINUED) CUCUMBERS, PEA, MUSTARD AND TURNIPS WITHIN 3 DAYS OF HARVEST.

COUSCOURS (CANTALOUPE, CUCUMBERS, MELONS, ...): FOR CONTROL OF APHIDS, SPIDER MITES, LEAFHOPPERS, GRASSHOPPERS, DIPTEROUS LEAF-MINER (ADULT), LYGUS BUGS, FALSE CHINCH BUG, CABBAGE LOOPER, SALT-MARSH CATERPILLAR AND CLIMBING OUTWORM APPLY 15-25 LBS. PER ACRE. DO NOT APPLY ON CANTALOUPE, CUCUMBERS, MELONS AND SUMMER SQUASH WITHIN 1 DAY OF HARVEST

FRUIT PLANTS, PEPPERS AND TOMATOES: FOR CONTROL OF APHIDS, SPIDER MITES, LEAFHOPPERS, AND GRASSHOPPERS APPLY 15-25 LBS. PER ACRE. DO NOT APPLY WITHIN 1 DAY OF HARVEST.

STRAWBERRIES: FOR CONTROL OF APHIDS, SPIDER MITES, LYGUS BUGS, GRASSHOPPERS, LEAFHOPPERS, LEAFROLLERS AND SALT-MARSH CATERPILLAR APPLY 30-50 LBS. PER ACRE. DO NOT APPLY WITHIN 1 DAY OF HARVEST.

U.S.D.A. Reg. No. 143-710

25-1969

CHEMICAL CO.

KEEP OUT OF REACH OF CHILDREN. POISONOUS IF SWALLOWED, INHALED, OR A SCRUBED TO SKIN. RAPIDLY ABSORBED THROUGH SKIN. DO NOT SWALLOW OR GET IN EYES, ON SKIN, OR ON CLOTHING. DO NOT BREATHE DUST. WASH THOROUGHLY WITH SOAP AND WATER AFTER HANDLING AND BEFORE EATING OR SLEEPING. WEAR CLEAN NATURAL RUBBER GLOVES, CLEAN PROTECTIVE CLOTHING AND GOGGLES. REPLACE GLOVES FREQUENTLY AND DESTROY USED GLOVES. WEAR A MASK OR RESPIRATOR OF A TYPE PASSED BY THE U. S. DEPARTMENT OF AGRICULTURE FOR PHOSDRIN INSECTICIDE PROTECTION. IF THE MATERIAL GETS INTO THE EYES, IMMEDIATELY FLUSH WITH WATER FOR AT LEAST 10 MINUTES AND GET MEDICAL ATTENTION. IN CASE OF SPILLAGE ON PERSON OR CLOTHING IMMEDIATELY REMOVE ALL CONTAMINATED CLOTHING AND WASH SKIN WITH SOAP AND WATER; WASH CLOTHING BEFORE RE-USE. DO NOT APPLY OR ALLOW TO DRIEF TO AREAS OCCUPIED BY UNPROTECTED HUMAN OR BENEFICIAL ANIMALS. UNPROTECTED HUMANS AND ANIMALS SHOULD BE KEPT OUT OF TREATED FIELDS OR DAYS OF TREATMENT. KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS. DO NOT CONTAMINATE FOOD OR FEED PRODUCTS. TO PROTECT FISH AND WILDLIFE, DO NOT CONTAMINATE STREAMS, LAKES OR PONDS WITH THIS MATERIAL.

DO NOT APPLY PHOSDRIN IF BEES ARE IN THE FIELD. TREAT ONLY LATE AT NIGHT AFTER BLOSSOMS CLOSE OR EARLY IN THE MORNING BEFORE BLOSSOMS OPEN.

BURY OR BURN EMPTY BAGS; NEVER RE-USE. WHEN BURNING, WORK TO WINDWARD TO AVOID DANGER FROM FUMES.

A PERMIT IS REQUIRED FOR APPLICATION OF PHOSDRIN IN CALIFORNIA. SEE YOUR LOCAL COUNTY AND AGRICULTURAL COMMISSIONER.

THIS PRODUCT WILL KILL FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. APPLY THIS PRODUCT ONLY AS SPECIFIED ON THIS LABEL.