

CASORON® G-10

NET CONTENTS

DICHLOBENIL WEED AND GRASS KILLER

50 POUNDS

8/16

CAUTION: KEEP OUT OF THE REACH OF CHILDREN

DIRECTIONS

NON-SELECTIVE WEED CONTROL

This product is designed primarily for industrial weed control. Annual and perennial broadleaf and grass weeds listed below are controlled in a wide range of non-cultivated situations. When applied as directed this product will give non-selective weed control for an extended period of time. Length of residual action is dependent upon rain-fall, soil type and season of application.

CASORON G-10 is a free-flowing granule, non-flammable and easy to use. Best results are obtained by distributing uniformly over the soil surface by hand operated or tractor mounted granule applicator.

CASORON G-10 is recommended for control of certain perennial weeds such as:

Artemisa	Fescue	Sheep Sorrel
Bindweed	Leafy Spurge	Timothy
Canada Thistle	Orchardgrass	Wild Artichoke
Curly Dock	Quackgrass	Wild Aster
Falsedandelion	Russian Knopweed	Wild Carrot
Catsear		Yellow Rocket

CASORON G-10 is recommended for control of annual grassy and broadleaved weeds and certain perennial weeds such as:

Annual	Foxtail	Purslane
Bluegrass	Gisela	Ragweed
Bluegrass	Groundsel	Red Deadnettle
Ball Thistle	Henbit	Redroot Pigweed
Camphorweed	Horsetail	Rosarypea
Carpetweed	Jerusalem oak	Russian Thistle
Chickweed	Goosefoot	Shepherdspurse
Citron Melon	Knotweed	Smartweed
Coffinweed	Lambsquarters	Spanishneedles
Crabgrass	Maypops	Spurge
Cudweed	Milkweed Vine	Teaweed
Dandelion	Morrena	Texas Panicum
Dogfennel	odorata	Hurrahgrass
Eveing	Miners lettuce	Wild Barley
prostrate	Natalgrass	Wild Mustard
Eddystone	Old Witchgrass	Wild Radish
Florida Purslane	Peppergrass	Yellow
Pringle	Pineappleweed	Woodsorrel
	Plantain	

For annual weed control CASORON G-10 should be applied prior to germination or when new plants are very small. For perennial weeds, make application during coldest weather when plants are fully dormant, or in the winter resting stage.

© 1965, Thompson Hayward, Inc. 11-72

ACTIVE INGREDIENT:

Dichlobenil* 10.0 %
INERT INGREDIENTS 90.0 %
 100.0 %

*2,6-dichlorobenzonitrile

A NON-SELECTIVE HERBICIDE FOR CONTROL OF WEEDS (BROADLEAF AND GRASSY) IN INDUSTRIAL AND NON-CROP AREAS.

CASORON is a Research Discovery of N. V. PHILIPS-DUPHAR U. S. Patent No. 3,027,248

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY.

PRECAUTIONS

This product is not selective in action and may destroy all types of vegetation. The treated area may be totally or partially non-productive for one or more years. Apply only to areas where complete plant control is desired. Do not apply to areas on slope above lawns or susceptible crops as run-off water from treated area could cause injury. Do not graze livestock in treated areas. Store in a tightly closed container in a dry place.

CAUTION

HARMFUL IF SWALLOWED. Do not breathe dust. Do not allow contact with eyes or on skin. Use with caution. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Do not store with propagative structures such as seed, bulbs, tubers, nursery stock, etc., or with feed or food products. Do not reuse container. Destroy when empty.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of the material when such use and or handling is contrary to label instructions.

The weeds listed above can be controlled in industrial and non-crop areas such as


- Electric substations
- Equipment sales and storage areas
- Lumber yards
- Around above ground pipes and tanks
- Non-surfaced roadways
- Railroad and highway right-of-way
- Petroleum installations
- Around buildings
- Along fence rows
- Farm fuel storage tanks and out buildings

For non-selective weed control CASORON G-10 should be applied at the rate of 120 to 200 pounds per acre (2 3/4 to 4 1/2 pounds per thousand square feet, 4 1/2 to 7 1/2 ounces per 100 square feet or 3 4 to 1 1/4 pounds per square rod). The lower rates may be used if the weed infestation is primarily annuals. For perennial weeds and in dry soil areas the higher rates should be used. Dead weeds from previous seasons growth do not need to be removed and soil should not be cultivated prior to application.

CASORON G-10 should be applied during the season of lowest temperatures. Apply November 1 to March 1 in most areas or until April 1 in the northern tier of states. Do not apply if air temperatures are expected to rise above 70°F within the following week. In areas of limited rainfall apply just prior to or during rainy season. For prompt and effective results rainfall or other precipitation should occur within a few days of application.

OTHER USES

UNDER ASPHALT: For general weed control under asphalt (in areas such as roadways, parking lots, recreational areas), after final grade is achieved, apply 100 to 120 lbs. per acre, (2.3 to 2.8 lbs. per 1,000 sq. ft.) Treated area should be covered with asphalt as soon as possible. For re-surfacing work, existing weeds should be sprayed with a contact herbicide prior to CASORON G-10 application.



Kansas City, Kansas 66110