

# DE-PESTER®

KELTHANE<sup>(R)</sup>, CAPTAN, SULFUR 3-10-10

## DIRECTIONS

**GRAPES:** Botrytis rot (bunch rot), Powdery mildew, silver mite, Brown almond (clover) mite, two-spotted mite, European red mite, Pacific mite, Willamette mite, and storage rots, 30-35 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically making up to 3 cover applications before bunches close. Do not apply within 7 days of harvest.

**POTATOES:** Early and late blight, two-spotted, Pacific and strawberry (Atlantic) mites apply 25-30 lbs. per acre. Begin applications when plants are 2 to 4 inches high and repeat at 7 to 10 day intervals as long as disease threatens.

**STRAWBERRIES:** Botrytis rot, powdery mildew, two-spotted, strawberry (Atlantic) and six-spotted mite, 30 lbs. per acre, immediately after picking. Repeat applications after picking as necessary. Do not apply within 2 days of harvest.

ACCEPTED

APR 16 1958

U.S. DEPARTMENT OF AGRICULTURE  
BUREAU OF PLANT INDUSTRY  
WASHINGTON, D.C.  
FOR ECONOMIC POISON REGISTER  
FILE UNDER NO. 148-743

**THOMPSON-HAYWARD CHEMICAL CO.**  
FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

## ACTIVE INGREDIENTS:

1, 1-bis(chlorophenyl)-2, 2, 2-trichloroethanol.....	3%
Captan(N-trichloromethyl Thioc-4-cyclohexene)	
1, 2-dicarboximide.....	10%
Sulfur.....	10%
INERT INGREDIENTS.....	77%
TOTAL.....	100%

(R) Kelthane is a trademark Reg. U. S. Pat. office For. & Haas Company

Directions (continued)

**TOMATOES:** (Established plants): Early blight, late blight, anthracnose, russet mite, two-spotted, Pacific and tropical mite. On young plants apply 25-30 lbs. per acre, increasing to 30-35 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5-7 day intervals as necessary for control. Because of the sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor. Do not apply within 2 days of harvest.

## CAUTION

KEEP OUT OF REACH OF CHILDREN!

Avoid inhalation of dust. Avoid prolonged or repeated contact with skin. Avoid contamination of feed and food-stuffs. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions.

The user is advised not to use sulfur on any crop unless local use has proved that sulfur is safe in that locality. STORE IN COOL, DRY PLACE. PROTECT FROM EXCESSIVE HEAT.

Burn bag immediately when empty. Stay out of smoke.

Read entire label. Use strictly in accordance with label cautions, warnings and directions, and in conformity with Federal and State regulations.

Highly toxic to fish. Do not contaminate any body of water nor apply to any area not specified on the label.

U. S. D. A. Reg. No. 148-743

NOTICE: No warranty, expressed or implied, is made concerning this product, its use, except that it conforms to the chemical description on this label and the person is cautioned to make any other representation. Neither Thompson-Hayward Chemical Company nor seller shall be liable for any damages resulting from handling, storage or use of this product.