

*resubmitted
9-21-67
BS*



COTTON DUST
5 - 40 - 0
SEVIN* Sulfur

CAUTION Keep Out of Reach of Children
SEE OTHER SIDE FOR ADDITIONAL CAUTIONS.

FOR THE CONTROL OF INSECTS ON COTTON

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-Methylcarbamate)	5% by wt.
Sulfur	40%
INERT INGREDIENTS55%
TOTAL	100%

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

ACCEPTED
NOV 13 1967

DEPESIER

COTTON DUST
5 - 40 - 0
SEVIN* Sulfur

CAUTION Keep Out of Reach of Children
SEE OTHER SIDE FOR ADDITIONAL CAUTIONS.

FOR THE CONTROL OF INSECTS ON COTTON

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-Methylcarbamate)	5% by wt.
Sulfur	40%
INERT INGREDIENTS	55%
TOTAL	100%

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

U. S. Pat. No. 2,903,478

USDA Reg. No. 148-

ACCEPTED

NOV 18 1957

U.S. DEPT. OF AGRICULTURE
 FOREIGN TRADE
 ADMINISTRATION
 REG. NO. 148-715
 TO ATTACHED CO.



THOMPSON-HAWKINS CHEMICAL COMPANY
Kansas City, Mo.

DIRECTIONS FOR USE

FOR EARLY-SEASON THRIPS, FLEA BEETLES AND COTTON FLEAHOPPER. ALSO FOR COTTON LEAFWORM AND STRIPED BLISTER BEETLE:

Apply 10 to 20 pounds per acre on appearance of insects.

FOR MID-AND LATE-SEASON BOLL WEEVIL, BOLLWORM, FALL ARMYWORM, COTTON LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS AND TARNISHED PLANT BUG. ALSO FOR LIGHT-TO-MODERATE INFESTATIONS OF LYGUS BUGS IN WESTERN IRRIGATED COTTON.

Apply 20 to 40 pounds per acre, depending on the size of the cotton and insect population level. Applications for boll weevil should be made every five to seven days as long as control is necessary.

FOR PINK BOLLWORM: Apply 30 to 50 pounds per acre on a five to seven day schedule depending on the size of the cotton and degree of infestation.

FOR STINK BUGS AND SALT-MARSH CATERPILLAR: Apply 40 pounds per acre as needed.

FOR GRASSHOPPERS: Apply 10 to 30 pounds per acre on appearance of insects and their feeding damage. The highest dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants. APHID populations will be suppressed by repeated applications of this insecticide.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

SEVIN may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

PEANUTS: For the control of Thrips, Leafhoppers, Three Corner Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetles, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Green Clover Worm-- apply 20 pounds per acre on appearances of insects and their feeding damage. Make application as necessary.

For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

HANDLING PRECAUTIONS

CAUTION. HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. Skin Contact With any Insecticide May be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note For Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atrophene is a tictol. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Do not contaminate or store near feed or food-stuffs.

Apply 10 to 20 pounds per acre on appearance of insects.

FOR MID-AND LATE-SEASON BOLL WEEVIL, BOLLWORM, FALL ARMY-WORM, COTTON LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS AND TARNISHED PLANT BUG. ALSO FOR LIGHT-TO-MODERATE INFESTATIONS OF LYGUS BUGS IN WESTERN IRRIGATED COTTON.

Apply 20 to 40 pounds per acre, depending on the size of the cotton and insect population level. Applications for boll weevil should be made every five to seven days as long as control is necessary.

FOR PINK BOLLWORM: Apply 30 to 50 pounds per acre on a five to seven day schedule depending on the size of the cotton and degree of infestation.

FOR STINK BUGS AND SALT-MARSH CATERPILLAR: Apply 40 pounds per acre as needed.

FOR GRASSHOPPERS: Apply 10 to 30 pounds per acre on appearance of insects and their feeding damage. The highest dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants. **APHID** populations will be suppressed by repeated applications of this insecticide.

SEVIN does not control spider mites. Where infestations are encountered, use a recommended miticide.

SEVIN may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

PEANUTS: For the control of Thrips, Leafhoppers, Three Corner Alfalfa Hopper, Cucumber Beetles, Bean Leaf Beetles, Japanese Beetle, Alfalfa Caterpillar, Velvet Bean Caterpillar, Green Clover Worm-- apply 20 pounds per acre on appearances of insects and their feeding damage. Make application as necessary.

For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

HANDLING PRECAUTIONS

CAUTION. HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do Not Take Internally. Avoid Prolonged or Repeated Breathing of Dust. Skin Contact With any Insecticide May be Harmful. Avoid Unnecessary Contact. Wash Hands Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily. Note For Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atrophene is antidotal. **KEEP OUT OF REACH OF CHILDREN AND ANIMALS. Do not** contaminate or store near feed or food-stuffs.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.