

# DE-PESTER<sup>TM</sup>

THIRAM, MANEB 3.3-6

### DIRECTIONS FOR USE

For application to either moist or dry foliage anytime dusting conditions are favorable. Application should begin when plants are small; increase dosage as plants grow larger. Dusters should be adjusted to give uniform and thorough coverage.

### RECOMMENDED USES

**TOMATOES:** For control of Botrytis gray mold, rhizoctonia fruit rot, early and late blight, anthracnose, stemphylium (gray leaf spot) and septoria leaf spots apply 20 - 30 lbs. per acre when disease is first reported in area and repeat at 7-10 day intervals, or oftener under severe late blight conditions.

### ACTIVE INGREDIENTS:

Thiram (Tetramethylthiuram disulfide).....	3.3%
Maneb (Manganous ethylene bisdithiocarbamate).....	6.0%
<b>INERT INGREDIENTS.....</b>	<b><u>90.7%</u></b>
<b>TOTAL</b>	<b>100.0%</b>

## CAUTION

**KEEP OUT OF REACH OF CHILDREN!**  
Harmful if swallowed. Avoid breathing of dust mist. Avoid contact with eyes skin, and clothing. Wash thoroughly after using. Avoid contamination of feed or foodstuffs. Keep away from children and domestic animals.

To protect fish and wildlife, do not contaminate streams, lakes or ponds with this material.

**THOMPSON-HAYWARD CHEMICAL CO.**  
FRESNO, CALIFORNIA — KANSAS CITY, MISSOURI

NOT A WARRANTY, EXPRESSED OR IMPLIED, IS MADE BY THIS COMPANY OR ITS AGENTS OR EMPLOYEES THAT IT CONFORMS TO THE CHEMICAL DESCRIPTION OF THIS PRODUCT AS SET FORTH OR IN ANY OTHER REPRESENTATION. Neither Thompson-Hayward Chemical Company nor seller shall be liable for any damages resulting from handling, storage or use of this product.