



SEVIN*-SULFUR 5-50 DUST

DIRECTIONS FOR USE

COTTON: For the control of early season thrips and cotton fleahopper, apply 10 to 20 lbs. per acre on appearance of insects. For control of mid and late-season boll weevil, bollworm, cotton leafworm, lygus and stink bugs and cotton leaf perforator, apply 20 to 40 lbs. per acre at from five to 7 day intervals on appearance of insects.

For salt-marsh caterpillar control: Apply 45 lbs. per acre.

For pink bollworm control on a 5 to 7 day achedule: Apply 30 to 45 lbs. per acre depending on the size of the cotton and degree of infestation. Sevin may not be applied after bolls open. Do not graze treated fields or feed trash to dairy or meat animals.

GRAPES: To control grape leafhopper and grape leaffolder, use 20 to 25 lbs. per acre when first brood appears. Use 20 to 40 lbs. per acre on second and third broods of grape leaffolder. May be applied on grapes up to harvest.

PEACHES, PLUMS, PRUNES: To control peach twig borer, use 40 to 50 lbs. per acre. May be used up to 1 day before peach harvest and within 3 days of harvest of plums and prunes.

ACTIVE INGREDIENTS:

| | |
|------------------------|-------|
| Carbaryl*..... | 5.0% |
| Sulfur..... | 50.0% |
| INERT INGREDIENTS..... | 45.0% |
| TOTAL...100.0% | |

*Trademark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

Directions for Use (continued)

Aphid population will be repressed when this insecticide is used in a regular schedule of treatments. Where mite infestations are encountered, use a recommended miticide. Sevin* is believed compatible with all pesticides commonly used on cotton.


CAUTION

KEEP OUT OF REACH OF CHILDREN! May be harmful if swallowed or inhaled. Avoid breathing dust. Avoid prolonged or repeated contact with skin. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling. Do not take internally.

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

U.S.D.A. Reg. No. 148-755



THOMPSON-HAYWARD CHEMICAL CO. FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

NOTICE: No warranty, expressed or implied, is made concerning this product or its use except that it conforms to the chemical description on this label and no person is authorized to make any other representation. Neither Thompson-Hayward Chemical Company nor seller shall be liable for any damages resulting from handling, storage or use of this product.