

RECOMMENDATIONS FOR USE

GREEN BEANS: For control of root, and stem rot caused by Rhizoctonia and Pythium, apply 20 lbs. per acre and distribute the dust over the seeds and surrounding soil as the seeds are planted.

COTTON: For control of damping-off caused by Rhizoctonia and Pythium, apply 50 pounds per acre and distribute over the seeds and surrounding soil as the seeds are planted.

ORNAMENTALS: For control of dampingoff caused by Rhizoctonia and Pythium, thoroughly mix 2 lbs. of dust per cubic yard of greenhouse soil.


To obtain satisfactory results use adequate dusting equipment and apply thoroughly with a uniform deposit of dust.

When used properly Captan will not harm seed, fruit, or foliage. Recommended applications leave no objectionable residues.

ACTIVE INGREDIENTS:

Pentachloronitrobenzene	10.0%
Captan (N∞frichloromethy)thi	io-
4-Cyclohexene-1, 2-	
dicarboximide)	10.0%
INERT INGREDIENTS	გი.ი%
TOTAL	100.0%

*Terraclor is a registered trademark of Olin Mathieson Chemical Corp.



10-2%

KEEP OUT OF REACH OF CHILDREN! Avoid contamination of feed and foodstuffs.

Avoid inhalation of dust or spray mist.

Avoid contact with skin. May cause skin irritation. Wash hands and face thoroughly after using.

Residues remaining of food crops at time of harvest should be removed by washing or by other effective means.

This product is toxic to fish. Keep out of lakes, streams, or ponds.

Apply this product only as specified on this label.

U.S.D.A. Reg. No. 148-780

THOMPSON-HAYWARD CHEMICAL CO.

FRESNO, CALIFORNIA

KANSAS

Y, KANSAS