

# DE-PESTER

THIRAM, SULFUR 7½-10

### RECOMMENDED USES

**STRAWBERRIES:** For control of Botrytis gray mold and for mildew apply 40 lbs. per acre and repeat at 10 day intervals beginning at early blossom stage. Make 3 to 5 applications. Good coverage of strawberry buds blossoms and developing fruit is necessary for best results. Remove residues by washing if applications are made within three days of harvest.

May cause some leaf burn in hot weather. Do not apply or allow to drift to sulfur sensitive crops.

### ACTIVE INGREDIENTS:

Thiram (tetramethyl-thiuramdisulfide).....	7.5%
Sulfur.....	10.0%
INERT INGREDIENTS....	82.5%
TOTAL.....	100.0%


# CAUTION

**KEEP OUT OF REACH OF CHILDREN!**

May cause irritation of eyes, nose, throat and skin.

Do not breathe dust. Harmful if inhaled or swallowed.

Do not get in eyes, on skin, or on clothing. Wash thoroughly with soap and water after handling; for eyes, flush with plenty of water and get medical attention. Consult local agricultural Authorities for further details as to time, frequency and rate of applications.



**THOMPSON-HAYWARD CHEMICAL CO.**  
FRESNO, CALIFORNIA — KANSAS CITY, MISSOURI

NOT A WARRANTY, expressed or implied, is made concerning this product, except as it conforms to the chemical description in this label and is not intended to make any other representation. Neither Thompson-Hayward Chemical Company nor seller shall be liable for any damages resulting from handling, storage or use of this product.