

NET CONTENTS:

MALATHION 5% DUST

POUNDS  
**ACCEPTED**  
**MAR 11, 1988**

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND RODENTICIDE ACT  
 FOR ECONOMIC POISON REGISTERED  
 UNDER NO. 148-793 SUBJECT  
 TO ATTACHED COMMENTS.

RECOMMENDED USES

Apply thorough, full-coverage application and repeat as needed.

**ALFALFA and CLOVER:** For control of aphids, spider mites, potato leafhopper, spittlebugs, grasshopper and alfalfa weevil larvae, apply 30 lbs. per acre. For control of spotted alfalfa aphid on Alfalfa apply 12-15 lbs. per acre by ground equipment or 9-16 lbs. per acre by airplane; use higher dosages on taller Alfalfa. Do not apply to Alfalfa in bloom. Do not apply within 7 days of harvest or pasturing.

**BLUEBERRIES:** For control of blueberry maggot apply 20 lbs. per acre. Do not apply within 1 day of harvest.

**COTTON:** For control of spider mites, aphids, leafhoppers, whiteflies, cotton leaf worm, leaf perforator and thrips apply 8-20 lbs. per acre.

**CRANBERRIES:** For control of leafhopper and black-headed fireworms apply 30-40 lbs. per acre by ground equipment. For aerial application use a minimum dosage of 50 lbs. per acre. Do not apply within 3 days of harvest.

**DATE:** For control of nitidulid beetles apply thoroughly to each cluster, not exceeding 40 lbs. per acre. Do not apply within 7 days of harvest.

**GRAZES:** For control of leafhoppers and spider mites apply 16-32 lbs. per acre and repeat as needed. Do not apply within 3 days of harvest.

**SQUASHES:** For control of aphids and spider mites apply 40 lbs. per acre. Do not apply within 3 days of harvest.

**WALNUTS:** For control of aphids and spider mites apply 32-48 lbs. per acre.

ACTIVE INGREDIENT :

Malathion (O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate).....

INERT INGREDIENTS.....

TOTAL..

5.0%  
 95.0%  
 100.0%

148-793

(Recommended Uses continued)

VEGETABLES:

**BEANS:** For control of Mexican Bean beetle and leafhoppers apply 30-35 lbs. per acre and repeat as needed. Do not apply within 1 day of harvest.

**CARROTS, and ENDIVE:** For control of aphids apply 24-28 lbs. per acre, except apply 40 lbs. per acre to Cauliflower. Do not apply within 7 days of harvest.

**CUCURBITS (Cucumbers, Melons, Pumpkins, and Squash):** For control of aphids, spider mites and leafhoppers apply 24-28 lbs. per acre. Do not apply unless plants are dry. Do not apply within 3 days of harvest.

**LETTUCE:** For control of aphids, spider mites, leafhoppers and cabbage looper apply 24-32 lbs. per acre. Do not apply within 7 days of harvest of head lettuce or within 14 days of leaf lettuce.


**MUSHROOMS:** For control of Phorid and Sciarid flies, apply 8-12 lbs. per acre or 1.4 lbs. per 60 ft. of single house as soon after picking as possible & repeat as necessary.

# Malathion 5% Dust

## CAUTION

KEEP OUT OF REACH OF CHILDREN!

Harmful if swallowed, Avoid breathing dust. Use a respirator. Avoid prolonged or repeated contact with skin. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Bury or burn empty bags; never reuse. When burning, work to windward to avoid danger from fumes. Consult local agricultural authorities for further details as to time, frequency and rate of application.



148-793

Do not apply Malathion if Bees are in the field. Treat only late at night after blossoms close, or early in the morning before blossoms open.

(Recommended Uses Continued)

MUSHROOMS (CONTINUED)

usually twice a week. Do not apply within one day of harvest.

ONIONS: For control of thrips apply 24-32 lbs. per acre. Do not apply within 3 days of harvest.

PEAS: For control of pea aphid apply 20 lbs. per acre. Do not apply within 3 days of harvest.

POTATOES: For control of aphids and leafhoppers apply 20-24 lbs. per acre.

SPINACH: For control of aphids apply 24-28 lbs. per acre. Do not apply within 7 days of harvest.

TOMATOES: For control of aphids, spider mites and tomato russet mite apply 24-36 lbs. per acre. Do not apply within 1 day of harvest.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do

empty bags; never reuse. When burning, work to windward to avoid danger from fumes. Consult local agricultural authorities for further details as to time, frequency and rate of application.

TO ATTACHED COMMENTS.  
148-793

Do not apply Malathion if Bees are in the field. Treat only late at night after blossoms close, or early in the morning before blossoms open.

(Recommended Uses Continued)

MUSHROOMS (CONTINUED)

usually twice a week. Do not apply within one day of harvest.

ONIONS: For control of thrips apply 24-32 lbs. per acre. Do not apply within 3 days of harvest.

PEAS: For control of pea aphid apply 20 lbs. per acre. Do not apply within 3 days of harvest.

POTATOES: For control of aphids and leafhoppers apply 20-24 lbs. per acre.

SPINACH: For control of aphids apply 24-28 lbs. per acre. Do not apply within 7 days of harvest.

TOMATOES: For control of aphids, spider mites and tomato russet mite apply 24-36 lbs. per acre. Do not apply within 1 day of harvest.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

THOMPSON-HAYWARD CHEMICAL CO.

FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

USDA Reg. No. 148-

Page 2 of 2