

MAR 26, 1969

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON CONTROL
ED UNDER NO.
TO APPROVED PRODUCTS

DIRECTIONS FOR USE

Measure the correct amount of Perthane E-4 and add to the sprayer tank while it is filling with the agitator running. Stir and mix thoroughly in hand sprayers. Thorough, full-coverage spray is important and nozzles should be adjusted properly for uniform distribution.

CHERRIES: Cherry fruit fly: Use 1 quart in 100 gallons of water. Thorough full-coverage applications should be made beginning when the adult flies appear. Repeat application every 10 days until harvest time. Do not apply within 2 days of harvest. An additional spray after harvest is suggested, especially if high temperatures have kept flies active.

PEARS: For pear psylla control use 1 to 2 quarts per 100 gallons of water. (8 quarts per acre). Apply as prebloom and or postbloom sprays and up to 2 cover sprays. Allow 30 days between cover sprays. Do not apply within 7 days of harvest.

This material may be used in orchards inter planted with apples, but the apples may not be used for apple pomace for livestock feed.

LETTUCE-Leafhoppers, imported cabbage worms, cabbage loopers and alfalfa loopers (California): Use 1 quart per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers. Apply when the insects are found to be present, making thorough application. Repeat at weekly to 10-day intervals or as required for adequate insect control. In hand sprayers use 2 teaspoons per gallon of water. Do not apply within 4 days of harvest.

SPINACH-Leafhoppers and cabbage loopers: Use 1 quart per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayers. Apply when insects are first noted, making thorough application. Repeat applications at weekly to 10-days intervals as required for adequate insect control. Do not apply within 7 days of harvest.

DE-PESTER

**PERTHANE
E-4**

(Contains 4 lbs. Perthane per gallon)

ACTIVE INGREDIENTS:	
Diethyl Diphenyl Dichloroethane	45.0%
Related Reaction products	2.3%
Xylene	48.7%
INERT INGREDIENTS	4.0%
TOTAL	100.0%

CAUTION: Keep Out Of Reach Of Children

(See Additional CAUTION Statement On Side Panel)



CAB
bage
lons
when
terva

This
Avoid
breast
dairy
forag

This
prode
equip

Do n

ESTER

**THANE
E-4**

(Perthane per gallon)

Perthane	45.00%
.....	2.30%
.....	48.70%
.....	4.00%
TOTAL	100.00%

Out Of Reach Of Children

(Additional CAUTION Statement On Side Panel)

T-H

DIRECTIONS FOR USE (CONT.)

CABBAGE, BROCCOLI CAULIFLOWER, BRUSSELS SPROUTS, KOHLRABI-Cabbage loopers and imported cabbage worms: Use 1 quart per acre in 25 to 100 gallons of water in ground sprayers or in 5 gallons of water in aerial sprayer. Apply when insects are first noted, making thorough applications at weekly to 10-day intervals as required for adequate insect control. Do not apply within 3 days of harvest.

CAUTION HARMFUL IF SWALLOWED

This solution if brought into contact with the skin may cause toxic symptoms. Avoid skin contact. In case of skin contact, wash with soap and water. Avoid breathing spray mist. Avoid contamination of feed or foodstuffs. Do not use in dairy barns, on dairy animals, animals being finished for slaughter or in feed or forage to be fed to these animals.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Do not reuse container. Destroy when empty.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

2M 1-69

USDA REG. NO. 148-805