

# DE-PESTER<sup>®</sup>

MANEB, SULFUR 6-25  
(AN AGRICULTURAL FUNGICIDE DUST)

## DIRECTIONS FOR USE

For application to either moist or dry foliage anytime dusting conditions are favorable. Application should begin when plants are small; increase dosage as plants grow larger. Dusters should be adjusted to give uniform and thorough coverage.

## RECOMMENDED USES

**BEANS:** (Dry Lima and String): For control of downy mildew, apply 15 to 35 lbs. per acre when mildew first appears and repeat at weekly intervals. Do not apply within 4 days of harvest.

**CARROTS:** For control of *Cercospora* leaf spot and *Alternaria* leaf blight, apply 15 to 45 lbs. per acre when diseases are reported in the area and repeat at 7 to 10 day intervals during the growing season, or as required for adequate control.

**CELERY:** For control of early and late blight, apply 15 to 45 lbs. per acre when plants first come through ground. Repeat at 3 to 5 day intervals until plants are set in field. Continue field applications on weekly schedule. Remove excess residues by stripping, trimming, washing or other effective means.

**EGGPLANTS, PEPPERS AND POTATOES:** For control of *Anthraco*se and *Cercospora* on Peppers; *Phomopsis*, early blight (*Alternaria*) and *Anthraco*se on Eggplant; and early blight (*Alternaria*) and late blight (*Phytophthora*) on Potatoes, apply 25 to 50 lbs. per acre. Begin application when plants are transplanted to the field or

## ACTIVE INGREDIENTS:

Maneb (Manganese Ethylene Bisdithiocarbamate).....	6.0%
Sulfur.....	25.0%
INERT INGREDIENTS.....	69.0%
TOTAL.....	100.0%

## RECOMMENDED USES (cont'd)

before diseases are expected to appear, and repeat at 7 to 10 day intervals or oftener under severe late blight conditions.

**ONIONS:** For control of downy mildew, *Botrytis* leaf blight and purple blotch, apply 25 to 50 lbs. per acre, beginning when plants reach the 4 leaf stage or when disease conditions are reported in the area and repeat at 7 to 10 day intervals during season.

**TOMATOES:** For control of early and late blight, *Anthraco*se, *Stemphylium* (Grey leaf spot) and *Septoria* leaf spots, apply 25 to 50 lbs. per acre when disease is first reported in area and repeat at 7 to 10 day intervals or oftener under severe late blight conditions.

**SPINACH:** For control of Downy Mildew (Blue Mold), apply 25 to 45 lbs. per acre at first signs of mildew and repeat at 7 to 10 day intervals or as required. Do not apply within 7 days of harvest and remove excess residues by washing.

# CAUTION

KEEP OUT OF REACH OF CHILDREN!

MAY CAUSE IRRITATION OF NOSE, THROAT, EYES AND SKIN. DO NOT BREATHE DUST.


In case of skin contact wash thoroughly with water.

STORE IN COOL, DRY PLACE. KEEP CONTAINER CLOSED WHEN NOT IN USE.

BURY OR BURN EMPTY BAGS: NEVER REUSE. WHEN BURNING WORK TO WINDWARD TO AVOID DANGER FROM FUMES.

Toxic to fish. Do not contaminate streams, lakes, or ponds.

USDA REG. NO. 148-310



## THOMPSON-HAYWARD CHEMICAL CO.

FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

NOTICE: This product is registered under the Federal Insecticide, Fungicide, and Rodenticide Act, and is subject to the provisions of that Act. It is unlawful to sell, distribute, or use this product in violation of the Act. For more information, contact the manufacturer or the appropriate state or federal agency.