

DIRECTIONS—BRUSH CONTROL
RIGHT OF WAYS • ROADSIDES
DRAINAGE DITCHES •

"Ammate" Weed and Brush Killer Solution, applied as a foliar spray, is highly effective for killing most woody plants, including hard-wood and coniferous species such as alder, ash, birch, cedar, elm, gum, hickory, maple, oak, pine, and willow.

FOLIAGE SPRAY APPLICATION

Prepare spray (water solution or oil water emulsion), and apply any time after brush has reached the full-leaf stage until foliage begins to discolor, preferably during periods of high temperatures and high humidity. Apply as a full-coverage spray to foliage, stems, limbs, and base of brush, thorough coverage is essential for best results. In roadside applications on tall, dense brush, it is often necessary to spray from field side as well as road side. Spraying away from crops planted close to right of ways will aid in preventing contact of crops with spray or drift; coarse sprays are less likely to drift. The product is not volatile.

SPRAY PREPARATION

Water Solution—For Each 100 Gals. of Spray: Use 10 gals. product with hydraulic equipment; use 18 to 72 gals. with air blast equipment. To improve wetting of foliage, add 4 to 6 fl. ozs. Du Pont Spreader-Sticker or 1 qt. Du Pont Surfactant WK.

Oil-Water Emulsion—For Each 100 Gals. of Spray: With hydraulic equipment, use 9 gals. product plus a pre-mix (prepared in water-free container) of 3/4 pt. Du Pont Emulsifying Agent A in 4 gals. of No. 2 fuel oil. Fill spray tank 3/4 full of water; start agitator, add product; add pre-mix; dilute to proper volume with water. For air blast equipment,

CONTINUED ON RIGHT PANEL

148-829

DE-PESTER[®]

ACCEPTED

OCT 13, 1971

UNDER THE FEDERAL INSECTICIDE ACT AND EXECUTIVE ORDER 11633 FOR ECONOMIC PESTICIDE REGISTRATION AND LABELING REQUIREMENTS TO ATTACHED DOCUMENTS

AMMATE*

WEED & BRUSH KILLER

SOLUTION

ACTIVE INGREDIENT	
Ammonium Sulfamate	48.4%
INERT INGREDIENTS	51.6%
10.9 Lbs./Gal. @ 60°F.	

CAUTION! MAY CAUSE SKIN IRRITATION.
 Avoid prolonged contact with skin.
 Keep Out of Reach of Children
 In case of contact, wash off with plenty of water.

GENERAL INFORMATION: This product is a water solution containing the equivalent of 5.6 lbs. Du Pont "Ammate" X Weed and Brush Killer per gal. It is highly effective for killing undesirable woody plants and for use as a contact spray for control of weeds and grasses. Temporary non-productivity of soil may be caused by heavy applications of the product; this condition usually disappears in the "over-winter" period.

Some species of woody plants and perennial weeds are difficult to control, and retreatment may be necessary if regrowth occurs.

This product is water-soluble, non-volatile, non-flammable,

It is non-selective; do not apply to, or

drain, or flush equipment on desirable plants or vegetation.

IMPORTANT: Wash sprayer thoroughly after use to remove all herbicidal solution and to reduce corrosion. Use of this product as an oil water emulsion minimizes corrosion of equipment. If oil water emulsion is not used, commercial equipment should be coated with an asphaltic base paint; or thoroughly wash exterior of equipment at end of each day and apply protective coating of oil each week or as often as practical. When equipment is down for week-end, wash exterior and coat with oil; for end-of-season storage, wash inside and outside and coat with oil.

USDA REG. NO. 148-829

*Registered trademark of E. I. du Pont de Nemours & Co. (Inc.)

55 U.S. GALS.

PACKED AND DISTRIBUTED BY



E. I. DU PONT DE NEMOURS & CO. (INC.)

WILMINGTON, DELAWARE

use 18 to 40 gals. product plus pre-mix of 1 qt. Du Pont Emulsifying Agent A in 4 gals. of No. 2 fuel oil; prepare spray mixture as above.

DIRECTIONS—WEED AND GRASS CONTROL

TENNIS COURTS • DRIVEWAYS
FENCE ROWS • INDUSTRIAL SITES

Applied as a water solution or as an oil-water emulsion, "Ammate" Weed and Brush Killer Solution is an effective contact spray for control of: **Woody Perennials** such as poison ivy, poison oak, and poison sumac; **Perennials** such as leafy spurge, bitter dock, goldenrod, perennial ragweed, milkweed, and blueweed; and **Annuals** such as crabgrass, broomweed, chickweed, cocklebur, Jimson weed, lambs'-quarters, larkspur, prickly lettuce, ragweed, and shepherd's-purse.

Use at rate of 18 gals. product per 100 gals. spray, either as water solution or as oil-emulsion. Prepare respective spray mixture as directed under "Brush Control" for hydraulic equipment. Thoroughly wet foliage and stems of undesired vegetation. The degree of control and duration of effect will vary with weed species, rainfall, temperature, and other conditions.

NOTICE: SELLER MAKES NO WARRANTY, EXPRESS OR IMPLIED, CONCERNING THE USE OF THIS PRODUCT. RISKY USES INDICATED IN THE LABEL. NEVER ASSESS ALL RISK OF USE AND/OR HAZARD OF THIS MATERIAL WHEN SUCH USE AND/OR HAZARD IS CONTRARY TO LABEL INSTRUCTIONS.

Manufactured by

E. I. DU PONT DE NEMOURS & CO. (INC.)

Industrial and Biochemicals Dept.

WILMINGTON, DELAWARE