

DIRECTIONS FOR USE

FRUIT AND NUT CROPS: Rates given in pints per 100 gallons unless specified. Aircraft — use equivalent rate in 10-50 gallons water per acre.

APPLES: 1 pint, red banded leafroller. Do not apply after second cover spray or within 40 days of harvest. Do not exceed 15 pints per acre.

CITRUS (GRAPEFRUIT, KUMQUATS, LEMONS, LIMES, ORANGES, TANGELOES, TANGERINES): 3-4 pints, grasshoppers, 1 pint, fire ants. Apply to trunk and main branches. Do not apply within 30 days of harvest. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not use over 10 pints per acre.

CRANBERRIES: 3 pints, grasshopper, 4 pints, armyworm. Do not apply after fruit begins to form.

PEARS: 1 pint, red banded leafroller, 1.2 pints, pear psylla. Do not apply within 40 days of harvest or after second cover spray. Do not exceed 15 pints per acre.

STRAWBERRIES: 2-3 pints per acre, cutworm (climbing only), strawberry crown borer, strawberry weevil. Do not apply after fruit begins to form.

FORAGE, FIELD, SEED AND VEGETABLE CROPS: Rates given in pints per acre. Do not exceed recommended rates. Ground rig use 15-100 gallons water per acre. Aircraft 10-50 gallons water per acre.

ALFALFA: Seed crop only. 2 1/2-3 1/2 pints, alfalfa looper, alfalfa weevil, armyworm, 1 1/2-3 1/2 pints, climbing cutworm, 3 pints, alfalfa caterpillar, pea aphids, webworm, 2 pints, clover leaf weevil, Lygus bug, grasshopper, 1 1/2 pints, spittle bug. Do not feed treated forage to dairy animals or animals being finished for slaughter.

BARLEY, OATS, WHEAT: 2 pints, armyworm, cutworms, grasshopper. Do not feed treated forage to dairy animals or animals being finished for slaughter. No restriction on the use of the grain.

BEANS (Dry, Green, Lima): 4 pints, armyworms, Mexican bean beetle, bean leaf beetle, cucumber beetle, blister beetle, flea beetle, cutworm, Lygus bug, bean leaf skeletonizer, garden webworm, pea weevil, 3 pints, corn earworm, grasshopper, serpentine leafminer, 1 pint, bean leafroller, bean leafhopper, 1.5 pints, thrips. No limitation on use of shelled beans as human food. Do not apply sprays to green or snap beans after pods begin to form. Do not feed treated vines to dairy animals or animals being finished for slaughter.

BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER: 4 pints, cabbage looper, corn earworm, diamondback moth, 3 pints, flea beetle, 2 pints, green peach aphid, serpentine leafminer, 1.5 pints, climbing cutworms, fall armyworm, thrips. Do not apply within 30 days of harvest or after edible parts start to form.

CABBAGE: 2 1/2 pints to control insects listed under Broccoli. Do not apply after heads start to form.

CARROTS: 3 pints, cutworm, flea beetle, 2 pints, grasshopper, 4 pints, Lygus bugs. No time limitation.

CELERY: 1.6 pints, cutworms, fall armyworms, green peach aphid, serpentine leafminer, thrips, 4-5 pints, celery leafhopper. Do not apply after celery plants start to bunch or after plants are half mature.

CORN: 2-6 pints, fall armyworm and corn earworm feeding in whorl, 3 pints, chinch bug, cutworm, flea beetle, grasshopper, 2 1/2 pints, webworm. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.

COTTON: 2-4 pints, armyworm, boll weevil, bollworm, cotton leafworm, cutworm (climbing), flea beetle, grasshopper, Lygus bug, thrips, 4 pints, stink bug, 3 pints, flea hopper. Do not graze dairy animals or animals being finished for slaughter in fields treated late in season.

EGGPLANT, PEPPERS: 2-3 pints, armyworm, blister beetle, Colorado potato beetle, cutworm, flea beetle, leafhopper, serpentine leafminer, 3 pints, cucumber beetle. Do not apply within 5 days of harvest. Do not apply after edible parts start to form.

LETTUCE: 4 pints, grasshopper, field cricket, 3 pints, cucumber beetle, flea beetle, 2 pints, armyworm, cutworm, 3-5 pints, cabbage looper. Do not apply after seedling stage-leaf lettuce. Do not apply after heads begin to form-head lettuce.

ONIONS: 2 pints, thrips. Do not apply to green or spring onions.

PARSNIPS: 1 1/2 pints, grasshopper, striped flea beetle. Do not use tops for feed or food.

RICE: 2 pints, armyworm, chinch bug, cutworm, grasshoppers, rice stink bug. Do not feed treated forage to dairy animals or animals being finished for slaughter. No limitation on use of grain.

RUTABAGAS: 2 pints, cutworm, 4 pints, flea beetle. Do not use treated tops for feed or food.

SOYBEANS: 2-4 pints, armyworm, bean leaf beetle, blister beetle, corn earworm, grass hopper, velvetbean caterpillar, 2 pints, cricket, 1 1/2 pints, flea beetles. Do not feed treated plants or ensilage made from treated plants to poultry, dairy animals or animals or animals being finished for slaughter. Do not feed soybean mill trash to livestock or poultry. Do not apply within 21 days of harvest.

SPINACH: 2 pints, armyworm, cutworm, green peach aphid, thrips, serpentine leafminer. Do not apply within 21 days of harvest. Do not apply more than once per season.

TOMATOES: 2-5 pints, armyworm, blister beetle, cabbage looper, Colorado potato beetle, cutworm, flea beetle, pinworm, russet mite, serpentine leafminer, tomato fruitworm, tomato hornworm, 3-5 pints, thrips, 4 pints, grasshoppers. Do not apply within 3 days of harvest.

WARNING

May be fatal if swallowed! Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after using.

Do not allow this product to drift onto adjoining food or feed crops as residues may contaminate these crops making them unfit for sale, feed or human consumption. Keep clear of food or dairy animals, streams, fish ponds, and wild life areas. Do not graze treated Cotton, or Feed gin waste to dairy animals or animals being finished for slaughter.

Do not store near feed or food products

This product is toxic to bees and should not be applied when bees are actively visiting the area.

This product is toxic to fish, shrimp, crab, birds and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

TOXAPHEN

(Contains 8 Lbs. Toxaphene)

ACTIVE INGREDIENTS:

Toxaphene*

Petroleum Distillate

INERT INGREDIENTS:

*Technical Chlorinated Camphene

WARNING: Keep Out

(SEE SIDE PANEL FOR ADDITIONAL INFORMATION)




APHENE E-8

8 Lbs Toxaphene Per Gallon


TS:	72%
late	20%
S:	8%
TOTAL	100%

ed Camphene (Chlorine content 67 - 69%)



Keep Out of Reach of Children
SEE PANEL FOR ADDITIONAL CAUTIONS

Reduce Hazards
**Follow This
 Rinse and Drain
 Procedure
 for
 Pesticide Containers**



Empty container into spray tank. Then drain in vertical position for 30 seconds

Add a measured amount of rinse water (or other diluent) so container is 1/4 to 1/5 full. For example, one quart in a one-gallon container



Rinse container thoroughly, pour into tank, and drain 30 sec. Repeat three times. Add enough fluid to bring tank up to level.

Crush pesticide container immediately. Sell as scrap for recycling or bury. Do not reuse.

