


Net Contents: Gallons



WARNING Keep Out of Reach of Children
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

Contains HPETACHLOR - 2 LBS. PER GALLON
ACTIVE INGREDIENTS:
Heptachlor*.....22.2%
Related Compounds..... 3.8%
Aromatic petroleum derivative
solvent.....04.0%
INERT INGREDINETS..... 50.0%
100.0%
*heptachlorotetrahydro-4,7-Methanoindene

This product is toxic to fish and wildlife. Kepp out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

Rinse spray equipment containers and dispose of liquid waste in a pit in non croplands located away from water supplies.

THOMPSON-HAYWARD CHEMICAL COMPANY
Subsidiary of Philips Electronics and Pharmaceutical Industries Corp.
Kansas City, Kansas 66110

USDA Reg. No. 148-

ACCEPTED
JUN 10 1968
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NOM 148-148 SUBJECT
TO ATTACHED COMMENTS.

DIRECTIONS

Before use read Directions Warning and Notice Statements Carefully

MIXING INSTRUCTIONS: Determine the amount of Heptachlor E-2 to use in the sprayer by multiplying the number of acres to be treated by the quantity of Heptachlor E-2 recommended per acre. Add this amount of Heptachlor E-2 to 1/2 the water required for coverage. Mix well then add remaining water. Continue mixing until spray mixture is uniform. If spray is allowed to stand, re-agitate before use.

For control of wireworms on Snap Beans, Lettuce, and Peppers, apply 1 gallon per acre in sufficient water to the soil prior to or at planting time. Work into the soil 4 to 6 inches.

ORNAMENTAL PLANTS & SHRUBS: For ants use 1 gallon in sufficient water per acre and work into the soil. For Asiatic Garden Beetle Larvae, European Chafer Larvae, Japanese Beetle Larvae, White-Fringed Beetle Larvae and White Grubs use 1 gallon in sufficient water per acre and wash into the top few inches of soil. For Chiggers, Fleas, and ticks, use 1/2 gallon in sufficient water per acre. Apply to grass, turf or ornamental plants. For wireworms use 1 gallon in sufficient water per acre, and work into the soil.

RUTABAGAS: For control of Root Maggots, use 1-1/2 - 2-1/2 gallons in sufficient water per acre. Work into top few inches of the soil prior to or at time of planting. Or use 1-1/2 gallons in sufficient water per acre as soon as seedlings are well up. Make a second application 3-4 weeks later. Do not use tops as food or feed.

WIREWORMS ON CORN: Mix 1 to 1-1/2 Gallons of Heptachlor E-2 in sufficient amount of water to cover one acre. Apply the spray to the ground before plowing and disc or work into the top 3 or 4 inches of soil. Corn may be treated with a row application of 1/2 to 3/4 gallons in sufficient water for one acre. prior to planting.

ROOT WORMS: For corn mix 1 to 2 quarts of Heptachlor E-2 in enough water to cover one acre. Apply 1 quart per acre as a row treatment. Apply 2 quarts per acre as a broadcast application. Apply the spray to the ground and disc or work into the top 3 or 4 inches of soil. If planter spray attachment is used, it may be applied as a band in the row at the time of planting corn.

WARNING

May be fatal if swallowed. Do not breathe vapor, dust or spray mist. Do not get on skin. In case of contact wash immediately with soap and water. Do not contaminate feed and foodstuffs. Misues as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTICE:

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and /or handling of this material when such use and/or handling is contrary to label instructions.

6-10-68
148-148
Comments