

DE-PESTER[®]

TEDION[®], DIBROM, SULFUR 2-4-20

DIRECTIONS FOR USE

Strawberries: For control of spider mites, aphids, and powdery mildew, apply 40 - 50 pounds per acre. Do not apply within 4 days of harvest, and do not apply within 35 day intervals during fruiting period.

CAUTION

This product is toxic to fish and wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment. Birds feeding on treated areas may be killed. Apply this product only as specified on this label.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

ACTIVE INGREDIENTS:

Tetradifon*	2.0%
Naled**	4.0%
Sulfur	20.0%
INERT INGREDIENTS...	74.0%
Total	100.0%

* 4-chlorophenyl 2, 4, 5-Trichlorophenyl Sulfone.

** 1, 2-dibromo-2, 2-dichloroethyl dimethyl phosphate.


® Tedion a research discovery of N.V. Phillips-Duphar, The Netherlands U.S. Patent No. 2,812,281

USDA REG. NO. 148-855

CAUTION

KEEP OUT OF REACH OF CHILDREN! Harmful if swallowed. May cause irritation of eyes, nose and throat. Avoid breathing dust. Avoid contact with skin and eyes. Wash thoroughly after using. Avoid contamination of feed and foodstuffs. Do not use on cherries as injury may result. May be toxic to bees, avoid application during periods of bee activity. May corrode metal dust equipment, clean equipment after use.

Sulfur may burn foliage when temperatures are excessively high.



BEST DOCUMENT AVAILABLE

THOMPSON-HAYWARD CHEMICAL CO.
FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.