



COPPER, SULFUR 15 - 50

DIRECTIONS FOR USE

Apply when air is calm at the first sign of infection or as recommended on the label for specific crops. When using ground equipment apply at the rate of 20 to 40 pounds per acre depending upon degree of infection and stage of plant development. Repeat at 7 - 10 day intervals as needed.

Before marketing food crops, remove residues by washing, stripping, wiping or other effective means.

RECOMMENDATIONS

GRAPES: Powdery Mildew (Sulfur can cause injury to foliage when applied just before or on days when the temperature exceeds 100°F.)

BEANS: Angular Leaf Spot, Anthracnose, Downy Mildew and Powdery Mildew. Apply as preventive dusting when plants are about 5 inches high. Repeat as necessary. Start with 20 pounds per acre and increase rate to compensate for plant development.

ACTIVE INGREDIENTS:

Copper Expressed as Metallic*.. 15%

Sulfur..... 50%

INERT INGREDIENTS..... 35%
100%

*Derived from Copper Oxide

CAUTION

KEEP OUT OF REACH OF CHILDREN!
Harmful if taken internally.

ACCEPTED
Sep 23/1968
NET WEIGHT 50 POUNDS
FURNISHED BY
FOR LOCATION
BY UNIT
TO ADDRESS

(RECOMMENDATIONS CONT'D)

EGGPLANT: Alternaria, Anthracnose, and Phomopsis.

CELERY, POTATOES: Early and late blight. On potatoes, make early preventive dustings at 20 to 30 pounds per acre.

PEPPER: Anthracnose, Cercospora, Leaf Spot and Smutty Mold.

SPINACH: Blue mold (downy mildew), white rust.

TOMATOES: Anthracnose, Early and Late Blight, Septoria Leaf Spot.

THOMPSON-HAYWARD CHEMICAL CO.
FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

NOTE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.