



THIO SULFUR 4-75


DIRECTIONS FOR USE

GRAPES: To control grape leafhopper and powdery mildew apply at 25 to 30 pounds per acre for early season control; apply at 30 to 35 pounds per acre for midsummer control. Do not use on concord variety of grapes as severe injury is likely to occur. For disease control, begin application when shoots are 6-18 inches long and dust every 14 days in each row middle or every 7 days in alternate middles.

Do not apply within 7 days of harvest.

TOMATOES & PEPPERS: Apply 25 lbs. per acre for the control of aphid, flea beetle, green stink bug, white fly, hornworm, and powdery mildew. Do not apply within 1 day of harvest.

Repeat at 7-10 day intervals to maintain control.



ACTIVE INGREDIENTS:

Endosulfan*.....4%
Sulfur.....75%

INERT INGREDIENTS.....21%
TOTAL.....100%

*Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide

CAUTION

Workers entering treated areas on the day of application should wear protective clothing. Use of this material in a manner or at a time other than in accordance with the directions may produce animal or plant injury, excessive residues or other undesirable results.

Do not apply during periods of extreme heat. Do not apply or allow to drift onto sulfur-sensitive varieties.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

USDA Reg. No. 148-872

WARNING

KEEP OUT OF REACH OF CHILDREN!

Hazardous if swallowed, inhaled or absorbed through skin. In case of contact, immediately remove contaminated clothing and flush skin or eyes with plenty of water; for eyes, get medical attention. Wash thoroughly with soap and water after handling and before eating or smoking. Wear clean clothing. During commercial or prolonged exposure in dusting operations, wear protective clothing including a mask or respirator of a type passed by the U.S. Department of Agriculture for Endosulfan protection. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

THOMPSON-HAYWARD CHEMICAL CO.

FRESNO, CALIFORNIA — KANSAS CITY, KANSAS

