

NET CONTENTS

GALLONS

[148-874]


# DE-PESTER®

# INSTITUTIONAL AREA SPRAY

### ACTIVE INGREDIENTS:

Petroleum Distillate . . . . .	98.92%
*Technical Piperonyl Butoxide . . . . .	.90%
Pyrethrins . . . . .	.18%
Total . . . . .	100.00%

\*Equivalent to 0.72% (butyl-carbityl) (6-propyl-piperonyl) ether and 0.18% related compounds.



**CAUTION** Keep Out of Reach of Children

### DIRECTIONS FOR USE

DE-PESTER Institutional Area Spray is a "Ready to Use" Insecticide to be used in appropriate spray equipment (fog applicators, sprayers, vaporizers or other dispensers) according to the following directions.

### FOOD PROCESSING PLANTS

Roaches, Waterbugs, Ants, Silverfish and Spiders: Apply as a coarse spray to all cracks and crevices where these insects normally hide. Spraying of dark places such as behind pipes, beneath sinks and along runways infested with roaches, waterbugs, ants, silverfish and spiders recommended. Hit these insects with spray whenever possible. Repeat application as often as required. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product as a space spray in these areas, apply only when plant is not in operation.

Bedbugs: Spray bed frames, springs, and all sides of mattresses, especially seams and tufts. Spray behind baseboards, loose wall paper and into all cracks around the room where this insect may hide and breed. Repeat application when necessary.

Clothes Moths and Carpet Beetles: Use as a fine spray. Apply lightly to clothing or interior surfaces of storage container such as closets or trunks. Repeat treatment as necessary. When spraying clothing for clothes moth control, pay particular attention to all seams and folds of garments. Repeat treatment every 30 days unless articles are stored in moth-tight containers. For Carpet beetles in food processing plants, treat same as roaches.

Flies, Mosquitoes and Gnats: Use as a mist spray. Close room and discharge spray throughout room. Use 1 ounce of spray for each 1000 cubic feet of space. Leave room closed for at least 10 minutes. Sweep up fallen insects. Repeat application when necessary. Avoid direct contact with food or food utensils.

STORED PRODUCTS IN ELEVATORS, GRAINAEIS, WAREHOUSES AND MILLS: For control of the exposed stages of Field Crickets, Granary Weevils, Confused Flour Beetles, Grain Moths, Cadelles, Rust Red Flour Beetles, Saw Tooth Grain Beetles, Spider Beetles, Grain Mites, Rice Weevils and Cheese Skippers: After thorough cleanup and prior to filling with grain apply at the rate of 1 gallon for each 1800 square feet of area. Wet surfaces of the storage area thoroughly. Hit insects directly with the spray whenever possible. Repeat application as often as necessary. Do not apply directly to grain or oil-bearing seeds.

### CAUTION

- KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.
- DO NOT SPRAY INTO OR NEAR OPEN FLAME.
- AVOID SPRAYING DIRECTLY ON FEED, FOODSTUFFS AND FOOD EQUIPMENT.
- KEEP CONTAINER TIGHTLY SEALED WHEN NOT IN USE.
- HARMFUL IF SWALLOWED.

### NOTICE

See the back of the container for complete instructions concerning the use of this product. The label indicated on the label. Buyer assumes all risk of use and contracting of this product. Do not use in a manner contrary to label instructions.

THOMPSON-HAYWARD CHEMICAL COMPANY  
Kansas City, Kansas 66110

USDA Reg. No. 148-874


NET CONTENTS

[148-874]

GALLONS

# DE-PESTER®


# INSTITUTIONAL AREA SPRAY



### ACTIVE INGREDIENTS:

Petroleum Distillate . . . . .	98.92 %
*Technical Piperonyl Butoxide . . . . .	.90 %
Pyrethrins . . . . .	.18 %
Total . . . . .	100.00 %

\*Equivalent to 0.72% (butyl-carbityl) (6-propyl-piperonyl) ether and 0.18% related compounds.



### CAUTION Keep Out of Reach of Children

### DIRECTIONS FOR USE

DE-PESTER Institutional Area Spray is a "Ready to Use" Insecticide to be used in appropriate spray equipment (fog applicators, sprayers, vaporizers or other dispensers) according to the following directions.

### FOOD PROCESSING PLANTS

Roaches, Waterbugs, Ants, Silverfish and Spiders: Apply as a coarse spray to all cracks and crevices where these insects normally hide. Spraying of dark places such as behind pipes, beneath sinks and along runways infested with roaches, waterbugs, ants, silverfish and spiders recommended. Hit these insects with spray whenever possible. Repeat application as often as required. Foods should be removed or covered during treatment. All food processing surfaces should be covered during treatment or thoroughly cleaned before using. When using this product as a space spray in these areas, apply only when plant is not in operation.

Bedbugs: Spray bed frames, springs, and all sides of mattresses, especially seams and tufts. Spray behind baseboards, loose wall paper and into all cracks around the room where this insect may hide and breed. Repeat application when necessary.

Clothes Moths and Carpet Beetles: Use as a fine spray. Apply lightly to clothing or interior surfaces of storage container such as closets or trunks. Repeat treatment as necessary. When spraying clothing for clothes moth control, pay particular attention to all seams and folds of garments. Repeat treatment every 30 days unless articles are stored in moth-tight containers. For Carpet beetles in food processing plants, treat same as roaches.

Flies, Mosquitoes and Gnats: Use as a mist spray. Close room and discharge spray throughout room. Use 1 ounce of spray for each 1000 cubic feet of space. Leave room closed for at least 10 minutes. Sweep up fallen insects. Repeat application when necessary. Avoid direct contact with food or food utensils.

STORED PRODUCTS IN ELEVATORS, GRAINARIES, WAREHOUSES AND MILLS: For control of the exposed stages of Field Crickets, Granary Weevils, Confused Flour Beetles, Grain Moths, Cadelles, Rust Red Flour Beetles, Saw Tooth Grain Beetles, Spider Beetles, Grain Mites, Rice Weevils and Cheese Skippers: After thorough cleanup and prior to filling with grain apply at the rate of 1 gallon for each 1800 square feet of area. Wet surfaces of the storage area thoroughly. Hit insects directly with the spray whenever possible. Repeat application as often as necessary. Do not apply directly to grain or oil-bearing seeds.

### CAUTION

- KEEP OUT OF REACH OF CHILDREN AND DOMESTIC ANIMALS.
- DO NOT SPRAY INTO OR NEAR OPEN FLAME.
- AVOID SPRAYING DIRECTLY ON FEED, FOODSTUFFS AND FOOD EQUIPMENT.
- KEEP CONTAINER TIGHTLY SEALED WHEN NOT IN USE.
- HARMFUL IF SWALLOWED.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than as indicated on the label. Buyer assumes all risk of use and handling of this product. Seller's liability is limited to the amount of purchase price. Use and handling contrary to label instructions.

THOMPSON-HAYWARD CHEMICAL COMPANY  
Kansas City, Kansas 66110