

around homes and
cessing plants, ware-
posed surfaces, such
of DE-FEND E-2 in

ns, litter and feed
k houses and milk

ble products areas of

contaminate water by

destroy by perforating

TRATED MATERIAL

mist. Use only with

case of contact with

minutes; get medical

minute food or feed

open flames.

as may be killed.

aged use, may cause

the use of this product

handling of this material



CAUTION Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.
DO NOT STORE BELOW FREEZING TEMPERATURES.

ACTIVE INGREDIENT:

Dimethoate (O, O-dimethyl S- [N-methylcarbamoylmethyl]
phosphorodithioate)* 23.50%

INERT INGREDIENTS 76.50%

100.00%

* (Contains 2.0 lbs. per gallon)

* Based on Dimethoate ROGOR® produced by Montecatini Edison

USDA Reg. No. 148-877

9-70



DIRECTIONS

**BEFORE USE READ DIRECTIONS, CAUTION AND
NOTICE STATEMENTS CAREFULLY**

HOUSEFLIES

RESIDUAL WALL SPRAYS: For the control of houseflies, including resistant strains, in dairy barns, hog pens, calf barns, poultry houses, and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint of DE-FEND E-2 in 3 gallons of water. Thoroughly wet all fly-resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 square feet of surface.

DE-FEND E-2 controls flies up to 8 weeks or longer.

Repeat applications should be made when necessary.

Remove dairy animals, calves under one month of age and poultry from building when applying residual wall sprays.

SPOT SPRAYS: For localized housefly control, apply a spray containing ½ pint of DE-FEND E-2 in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways, around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

MAGGOT SPRAYS: For the control of housefly maggots, mix ½ pint DE-FEND E-2 in 5 quarts of water and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged-layer houses, garbage dumps and manure piles.

Repeat application as additional manure or garbage is added.

DIRECTIONS--Continued

CAUTION

NOTICE



DE-FEND™ E-200

DIMETHOATE SYSTEMIC INSECTICIDE

CAUTION: Keep Out of Reach of Children

ACTIVE INGREDIENT

INERT INGREDIENTS



DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

HOUSEFLIES

RESIDUAL WALL SPRAYS: For the control of houseflies, including resistant strains, in dairy barns, hog pens, calf barns, poultry houses, and other farm buildings, apply a 1% residual spray to the ceilings, walls, stanchions, etc. Prepare the spray by mixing 1 pint of DE-FEND E-2 in 3 gallons of water. Thoroughly wet all fly-resting areas to the point of run-off. One gallon of spray will cover 500 to 1,000 square feet of surface.

DE-FEND E-2 controls flies up to 8 weeks or longer.

Repeat applications should be made when necessary.

Remove dairy animals, calves under one month of age and poultry from building when applying residual wall sprays.

SPOT SPRAYS: For localized housefly control, apply a spray containing 1 pint of DE-FEND E-2 in 5 quarts of water with a knapsack or similar type sprayer to areas frequented by flies, such as doorways around windows, etc.

Repeat applications should be made when necessary. Good sanitation is a necessary part of any effective fly control program.

MAGGOT SPRAYS: For the control of housefly maggots, mix 1/2 pint DE-FEND E-2 in 5 quarts of water and apply as a coarse spray or with a sprinkling can to fly-breeding areas, such as poultry droppings in caged-layer houses, garbage dumps and manure piles.

Repeat application as additional manure or garbage is added.