
DIRECTIONS FOR USE

GRAPES: To control spider mites, and leafhoppers, use 20 to 30 LBS. per acre. Do not apply more than twice during fruiting season or within 15 days of harvest.

COTTON: To control spider mites, stink bugs, and lygus bugs use 30 to 35 LBS. per acre. Do not graze treated fields or feed treated plants to meat/dairy animals. Do not apply after bolls open.

Do not apply when weather conditions favor drift from areas treated.

Apply this product only as specified on this label.




Active Ingredients

Ethion*......4.0 Endosufan**.....3.0 Inert Ingredients...§ 3.0

*0,0,0', 0'-Tetraethyl S,S'-methylene bisphosphoro-dithioate.

**Hexachlorohexahydro methano-2,4,3-benzodioxathiepin oxide.



MEED OUT OF REACH OF CHILDREN!

Avoid breathing dust. May be fatal if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and gog- -. gles. In case of contact, wash immediately with soap and water. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contominate feed and food stuffs. Wash hands, arms, and face thoroughly with soap and water before eating and smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish & wildlife. Keep out of lakes, streams, or ponds. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse container. Destroy when empty.

USUA Reg. -40. 148-819