



DIRECTIONS FOR USE

FOR EARLY-SEASON THRIPS, FLEA BEETLES AND COTTON FLEAHOPPER. ALSO FOR COTTON LEAFWORM AND STRIPED BLISTER BEETLE:

Apply 5 to 10 pounds per acre on appearance of insects.

FOR MID- AND LATE-SEASON BOLL WEEVIL, BOLLWORM, FALL ARMYWORM, COTTON LEAF PERFORATOR, LEAF ROLLERS, LEAFHOPPERS AND TARNISHED PLANT BUG. ALSO FOR LIGHT-TO-MODERATE INFESTATIONS OF LYGUS BUGS IN WESTERN IRRIGATED COTTON.

Apply 10 to 20 pounds per acre, depending on the size of the cotton and insect population level. Applications for boll weevil should be made every five to seven days as long as control is necessary.

FOR PINK BOLLWORM: Apply 15 to 25 pounds per acre on a five to seven day schedule depending on the size of the cotton and degree of infestation.

FOR STINK BUGS AND SALT-MARSH CATERPILLAR: Apply 20 pounds per acre as needed.

FOR GRASSHOPPERS: Apply 5 to 15 pounds per acre on appearance of insects and their feeding damage. The highest dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants.

APHID populations will be suppressed by repeated applications of this insecticide.

SEVIN does not control spider mites. Where infestations are encountered use a recommended miticide.

SEVIN may be applied after bolls open. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

For protection of honeybees avoid use, if possible, during periods when honeybees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to treat hives at a safe distance until one week after application.

HANDLING PRECAUTIONS

CAUTION. HARMFUL IF INHALED OR SWALLOWED. Do not breathe Dust. Do not take internally. Avoid prolonged breathing of Dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIANS. Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidote. KEEP OUT OF REACH OF CHILDREN AND ANIMALS. DO NOT

DE-PESTER®

COTTON DUST
10 — 75 — 0
SEVIN* SULPHUR

ACCEPTED
MAY 11, 1971
UNDER THE FEDERAL INSECTICIDE
ACT OF 1947 AND UNDER THE
FEDERAL TOXIC SUBSTANCE CONTROL
ACT OF 1972. THIS PRODUCT IS
REGISTERED UNDER THE FEDERAL
INSECTICIDE ACT OF 1947. SEE
ATTACHED COMMENTS

CAUTION Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-Methylcarbamate) 10%
Sulphur (93% thru 325 mesh) 75%

INERT INGREDIENTS

15%
100%

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

U. S. Pat. No. 2,903,478