


NET CONTENTS

GALLON



HEPTACHLOR E-3


3 LBS. PER GALLON

WARNING - Keep Out of Reach of Children

ACTIVE INGREDIENTS:

Heptachlor*	31.7%
Related Compounds	11.7%
Xylene	51.6%
INERT INGREDIENTS	<u>5.0%</u>
	100.0%

*(Heptachlorotetrahydro-4,7-methanoindene)



This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

ANTIDOTE

EXTERNAL — In case of spillage on skin, wash promptly with soap and water.
INTERNAL — Emetic of 1 tablespoonful mustard in tumbler of water. CALL A PHYSICIAN.

WARNING

May be fatal if swallowed. Do not breathe vapor, dust or spray. Do not get on skin. In case of contact wash immediately with soap and water. Do not contaminate feed and food-stuffs. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops or cause residues in excess of official tolerances.

Do not reuse container. Destroy when empty. Not for use or storage in or around the home. Do not use, pour, spill or store near heat or open flame.

DIRECTIONS

BEFORE USE READ DIRECTIONS, WARNING AND NOTICE STATEMENTS CAREFULLY.

MIXING INSTRUCTIONS: Determine the amount of Heptachlor E-3 to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Heptachlor E-3 recommended per acre. Add this amount to 1/2 the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. If spray is allowed to stand, re-agitate before use.

ROOT WORM: For Corn use 1 1/2 to 2 1/2 pints per acre. Apply the spray to the ground and disc or work into the top 3 or 4 inches of soil. If planter spray attachment is used, it may be applied as a band in the row at the time of planting corn.

WIREWORMS ON CORN: Use 5 pints to 1 gallon per acre. Apply the spray to the ground before plowing and disc or work into the top 3 or 4 inches of soil. Corn may be treated with a row application of 2 1/2 pints to 1/2 gallon per acre, prior to planting.

NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and or handling of this material when such use and or handling is contrary to label instructions.