

Mar 7, 1973

148-981

### DIRECTIONS—Continued

**SMALL GRAINS—(BARLEY, OATS, RYE AND WHEAT) ARMY CUTWORM, PALE WESTERN CUTWORM**—Apply 1 to 1 1/4 pints per acre each application in sufficient water for good coverage. Do not use after heads begin to form. Do not pasture animals in treated fields. Make only a single application, do not apply within 45 days of harvest.

Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

Use this product only as stated on the label.

**DANGER:** Post treated areas. Do not treat areas where runoff will contaminate water used


for domestic purposes or contaminate streams, lakes, or ponds. Do not graze orchards, cut for hay or allow livestock to clean up drops

**DANGER:** Poisonous by Skin Contact, Inhalation or Swallowing! Rapidly Absorbed Through Skin! Do not get in eyes, on skin or on clothing. Wear a mask or respirator of a type passed by the U. S. Bureau of Mines for Endrin protection. In case of contact, immediately remove all contaminated clothing and flush skin or eyes with plenty of water for at least 15 minutes; for eyes, get medical attention. Do not breathe spray mist. Wash thoroughly with soap and water after handling and before eating or smoking; wear clean clothing. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals. Misuse as to quantity, timing or method of application can cause damage or injury to animals, persons, property or crops. Wear clean synthetic rubber gloves

### DESTROY CONTAINER WHEN EMPTY

Flush out thoroughly with water in some location where flush water can be run off harmlessly. Then punch container full of holes or crush so that it can not be used again for any purpose. Take to city dump or some other type location where no attempt will be made to salvage for any use whatsoever.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and handling of this material when such use and/or handling is contrary to label instructions.



# ENDRIN E-1.6

1.6 Lbs. Per Gallon



**DANGER** Keep Out of Reach of Children



SEE ANTIDOTE STATEMENT AND OTHER REQUIRED WARNING STATEMENTS ON OTHER PANELS.

## POISON

### ACTIVE INGREDIENTS:

Endrin\* ..... 19.5%

Xylene ..... 76.5%

INERT INGREDIENTS ..... 4.0%

100.0%

\*(Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene)

EPA Reg. No. 148-981

SG-1-73-A



THOMPSON-HAYWARD  
CHEMICAL COMPANY

P.O. BOX 2383 KANSAS CITY, KANSAS 64110

BEF  
MIXING IN  
of Endrin E  
multiplying  
by the qua  
per acre. A  
1/2 the wat  
age Mix w  
tinue mixing  
spray is allo

For ground  
of water pe  
from 3-10

**NOTE:** A  
weather. In  
until redisso

**COTTON-BO**  
bollworm):

**INTE**  
glass  
**EXTE**

**NOTE**  
antic  
be ne



# ENDRIN E-1.6

1.6 Lbs. Per Gallon



**DANGER** Keep Out of Reach of Children



SEE ANTIDOTE STATEMENT AND OTHER REQUIRED  
WARNING STATEMENTS ON OTHER LABELS

**POISON**

**ACTIVE INGREDIENTS:**

Endrin\* ..... 19.5%

Xylene ..... 76.5%

**INERT INGREDIENTS** ..... 4.0%

100.0%

\*(Hexachloroepoxyoctahydro-endo, endo-dimethanonaphthalene)

EPA Reg. No. 148-981

SG-1-73-A



**THOMPSON HAYWARD  
CHEMICAL COMPANY**

P.O. BOX 2183 KANSAS CITY, KANSAS 64110

## DIRECTIONS

BEFORE USE READ WARNING, CAUTION AND NOTICE STATEMENTS CAREFULLY

**MIXING INSTRUCTIONS:** Determine the amount of Endrin E-1.6 to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Endrin E-1.6 recommended per acre. Add this amount of Endrin E-1.6 to 1/2 the water required for complete, full coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. If spray is allowed to stand, reagituate before use.

For ground application use from 5-15 gallons of water per acre. For aerial application, use from 3-10 gallons per acre.

**NOTE:** A precipitate may deposit in cold weather. In such case, warm gently and agitate until redissolved before using.

**COTTON-BOLL WEEVIL, BOLLWORM** (not pink bollworm): Apply one pint per acre for early

season control. For midseason or late season control use 1 to 2 pints per acre. Spray on 5 to 8-day schedule until control is obtained. Resume treatment if insects reappear. **FLEA-HOPPERS, THRIPS:** Use 1/2 pint per acre per application for early season control. For mid-season control use 3/4 pint and for late season control use 1 pint per acre. Repeat treatments as necessary for continued control. **COTTON LEAFWORM, LEAF PERFORATOR, LYGUS BUGS, TARNISHED PLANT BUGS AND SALT-MARSH CATERPILLAR, CUTWORMS—**Use 1 to 2 pints per acre each application, when insects appear. Apply to give uniform coverage. Repeat as necessary. Workers entering treated cotton fields within 5 days of treatment should be protected. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not graze dairy animals or animals being finished for slaughter on treated fields, or be fed gin waste.

### ANTIDOTE

**INTERNAL:** If any of this insecticide is swallowed, give a tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. Keep victim prone and quiet.

**EXTERNAL:** If skin is contacted, wash immediately with soap and water.

### CALL A PHYSICIAN IMMEDIATELY

**NOTE TO PHYSICIAN:** In case of poisoning, administer barbiturates as for anticonvulsant therapy. Observe patient carefully since repeated treatment may be needed.