



READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNING AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.



DIRECTIONS

Use 2 level tablespoonfuls SEVIN W-50 in 1 gallon of water. To use, mix designated amount of SEVIN W-50 in a little water to make a thin paste, then add required amount of water. Do not store diluted spray. Make new dilution for each use.

LAWN INSECTS: Chinch Bugs, Sod Webworms, Earwigs, Leafhoppers, Millipedes, Mosquitoes, Fall Armyworms, Fleas and Ants — Use 6 ozs. in 15 gallons of water and cover 750 sq. ft. of established lawn. Mow lawn first and apply after rain or watering. Do not water for at least two days after application. Repeat in 2 or 3 weeks or as necessary. Keep children and pets off treated area until this material is washed into the soil and the area is completely dry.

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Japanese Beetles, Aphids, Bagworm, Elm Leaf Beetle, Certain Leafminers, Rose Slug, Scale Insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, Exposed Thrips and certain other insects — Start spraying when insects or their damage appear. Repeat weekly or as necessary. Do not use on Boston Ivy.

FRUIT TREES: APPLES, PEARS AND PEACHES: Apple Aphid, Leaf Rollers, Lesser Peach Tree Borer, — Make one application between bud break and bloom, a second application at petal fall and then at 3 to 4-week intervals for 3 to 4 months as necessary. Rosy Apple Aphid — Apply before leaves are curled. May be applied within 1 day of harvest. Apple Maggot, Codling Moth, Oriental Fruit Moth and Plum Curculio — Apply at petal fall, and every 10 to 14 days thereafter until control is achieved.

GRAPES: Use 2 to 4 pounds for European fruit lecanium, grape leaf folder, grape leafhopper and grape leaf skeletonizer. Apply just before first brood leaf folder larvas emerge from rolls and as needed for leafhoppers. Use 4 pounds for grape berry moth, Japanese beetle, June beetles and red-banded leaf roller. A dilute spray of 200 gallons per acre is suggested.

VEGETABLES: POTATO, EGGPLANT, PEPPERS, CUCUMBERS, SUMMER SQUASH: Use 1 to 2 pounds for Colorado potato beetle, flea beetles and leafhoppers. Use 2 to 4 pounds for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. On potato, tomato, and eggplant, use 4 pounds for cutworm control. Where aphids are a problem on potatoes, use one pound of DE-FEND W-25. (1/4 pound dimethoate per acre.)

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of dust or spray mist. Avoid contact with skin; wash thoroughly after using. Avoid storage near feed and food products. Keep out of reach of children. To avoid possible fruit thinning on apples and foliar injury on McIntosh do not apply within 15 days after petal fall. Dispose of container promptly when empty.

NOTICE

ACC
1973
148-750

SEVIN* W-50

CAUTION Keep Out of Reach of Children

SEE BACK PANEL FOR ADDITIONAL CAUTIONS.

KILLS SOD WEBWORMS—BAG WORMS AND CERTAIN INSECT PESTS OF LAWNS AND ORNAMENTALS

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 50.0%

INERT INGREDIENTS 50.0%

100.0%

*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate
U.S. Pat. No. 2,903,478