

### DIRECTIONS—Continued


residue. After spraying the inside of bins, allow at least 24 hours for the spray to dry before filling bin with grain. Spray outside of the bins every four to five weeks until three applications have been made. Then spray outside of bins once every three months. One gallon of Farm Bin Spray treat from 500 to 1,000 square feet, depending on surface absorption.

If grain is already infested when placed in the farm bins, fumigate with grain fumigants. Insects crawling out of fumigated grain and contacting sprayed bin surfaces, should be killed by Farm Bin Spray residue.

### CAUTION

Do not apply directly to food products. Avoid inhalation of spray mist or contact with skin or eyes. DO NOT POUR, SPRAY OR STORE NEAR FIRE OR FLAME. Harmful if swallowed. Keep out of reach of children. Do not reuse empty container. Perforate or crush and bury with waste.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.



# FARM BIN SPRAY

Contains METHOXYCHLOR

### CAUTION Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.


#### ACTIVE INGREDIENTS:

Technical Methoxychlor*	5.00%
Xylene	30.00%
Aliphatic Petroleum Derivative Solvent	65.00%
	100.00%

\*Equivalent to 4.4% 2,2-bis (p-methoxyphenyl)-1,1,1-trichloroethane and 0.6% other isomers and related compounds.

EPA Reg. No. 148-994

5-72



THOMPSON-HAYWARD  
CHEMICAL COMPANY

P O BOX 2383 KANSAS CITY, KANSAS 66110

BEFORE USE

Kills on  
Lesser G  
Toothed

FARM B  
obtain be  
and area  
attacking  
following

1. Clean  
where

2. Clean  
debris

3. Spray  
to we

4. Spray  
the w

A coarse  
penetrat