

COPPER - SULFUR DUST FOR PEANUTS

FOR THE CONTROL OF LEAF SPOT AND LEAF HOPPER

For the Bunch or Runner types make first application during the first ten days of July. On Spanish, dusting should be started about ten days earlier following with two or three applications at as nearly fourteen day intervals as weather conditions will permit.

The rate of application per acre should range from 15 to 20 lbs. per acre totaling approximately 70 lbs. per acre in four applications or 50 lbs. in three.

Should an application be washed off within a few hours, the treatment should be repeated as soon as weather conditions will permit.

ACTIVE INGREDIENTS:

Copper (in copper oxide form) expressed as metallic	4.00%
Sulfur	75.00%
INERT INGREDIENTS:	21.00%
TOTAL	100.00%

CAUTION KEEP OUT OF REACH OF CHILDREN

EPA REG. No. 148-1004

THOMPSON-HAYWARD CHEMICAL CO.

PLANTERS CHEMICAL DIVISION

NORFOLK, VA.

FAYETTEVILLE, N. C.

ALBANY, GA.

CAUTION

Harmful if swallowed. Sulfur may burn foliage when temperatures are excessively high (90° F. or above).

Do not reuse container. Destroy by burying in a safe place away from water supplies. Do not burn.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

