



20% TOXAPHENE DUST

CABBAGE, BROCCOLI, LETTUCE AND BRUSSELS SPROUTS: To control Cabbage Loopers and other Caterpillars - apply at the rate of 10 lbs. per acre. For Cutworms (soil) - apply 20 lbs. per acre to soil surface. For Cabbage and Head Lettuce, apply up to 7 days before harvest, if the outer leaves are stripped at harvest. If the outer leaves are to be left on at harvest, do not apply after the heads begin to form. For Leaf Lettuce, do not apply after seeding is over. For Broccoli and Brussels Sprouts, do not apply after the edible parts begin to form. On all the above, do not apply this product if leaves, stems, or heads are to be fed to dairy or meat animals.

COLLARDS AND KALE: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface-Feeding Cutworms - Apply at the rate of 10 to 20 lbs. per acre. Do not apply more than 4 times nor more than 3 times after thinning. Do not apply within 14 days of harvest when 10 lbs. per acre is used; do not apply within 21 days of harvest when 20 lbs. per acre is used.

SPINACH: To control Army Worm, Cabbage Worm, Cabbage Looper, Climbing and Surface-Feeding Cutworms - Apply only 1 time at the rate of 10 lbs. per acre. Do not apply within 21 days of harvest.
BLANS: To control Cutworms (soil) - apply 10 lbs. per acre on soil in late afternoon. Do not apply after buds begin to form.

TOMATOES: To control Tomato Hornworm, Tobacco Fruit Worm, Spiny-tipped Leaf Miner, Flea Beetle, Cutworms, Cabbage Looper, Russet Mites and Colorado Potato Beetles - apply 10 to 20 lbs. per acre as required. Do not apply within 5 days of harvest.

SOYBEANS: To control Soybean Bean, Cutworm, Green Earworm, Bean Leaf Beetle, Army Worms and related species - apply at the rate of 20 lbs. per acre. Do not apply within 21 days of harvest. Do not apply to plants that are to be used for animal feed. Do not apply to plants that are to be used for dairy animals or meat animals.

PEANUTS: To control Fall Army Worm, Green Earworm, and related species - apply at the rate of 20 lbs. per acre. Do not apply within 21 days of harvest.

CORN AND SORGHUM: To control Spring and Fall Army Worm - apply at the rate of 10 lbs. per acre. On Sorghum, do not apply after harvest, nor more than once a year. Do not graze dairy animals on lush or treated forage. Do not use on treated forage.

WHEAT, BARLEY, OATS AND RYE: To control Climbing and surface-feeding Cutworms, Grasshoppers, and Chinch Bugs - apply at the rate of 10 lbs. per acre. Do not apply more than once after heads have begun to form. Apply within one week of harvest for Wheat and Oats; within 2 weeks of harvest for Rye. Do not graze dairy animals or meat animals on treated fields.

CAUTION

Harmful if swallowed. Avoid pro-
of dust mist. Avoid contact with
clothing. Wash thoroughly. Avoid
food or foodstuffs. Keep away from
domestic animals. Do not use on
use on feed or forage.

This product will kill fish and will
any body of water.

Do not prevent residue completely
dose of waste pesticide by burning
immediately and stay out of stacks. I
Seller warrants that the product
chemical description and its reason-
purpose stated on the label when
used with directions under warranty
out neither this warranty nor any
of MERCHANTABILITY or FITNESS FOR
purpose, express or implied, extend
to the product contrary to the label
if the abnormal conditions are not
caused by the use of the product or
the conditions of the environment.

ACTIVE INGREDIENTS

Toxaphene-Technical Chlorinated Camphene 20.00
(Chlorine Content 67-69%)

INERT INGREDIENTS

80.00
100.00

CAUTION KEEP OUT OF REACH OF CHILDREN.

USDA Reg. No. 1863-64

P-C

PLANTERS CHEMICAL COMPANY

Division of The Georgia Haywood Chemical Company

NORFOLK, VA FAYETTEVILLE, N. C. ALBANY, GA

ACCEPTED
 MAY 9, 1968
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC PEST CONTROL SUBJECT
 REGISTERED UNDER NO. 1868-64
 TO ATTACHED COMPLETES

148-1014

20% TOXAPHENE DUST

ACTIVE INGREDIENTS

Toxaphene-Technical Chlorinated Camphene 20.00
 (Chlorine Content 67-69%)

INERT INGREDIENTS

80.00
 100.00

CAUTION KEEP OUT OF REACH OF CHILDREN.

U.S. EPA REG. NO. 1868-64

P-C

PLANTERS CHEMICAL COMPANY

Division of Thompson Hayward Chemical Company

NORFOLK, VA. FAYETTEVILLE, N. C. ALBANY, GA.

CORN AND SORGHUM: To control Army Worm and Fall Army Worm — apply at the rate of 15 lbs. per acre. On Sorghum, do not apply within 40 days of harvest, not more than once after heads start to form. Do not graze dairy animals or animals being finished for slaughter on treated fields. Do not ensile treated forage.

WHEAT, BARLEY, OATS AND RYE: To control climbing and surface feeding Cutworms, Army Worms, Grasshoppers, and Chinch Bugs - - apply at the rate of 10 lbs. per acre. Do not apply more than once after heads have begun to form. Do not apply within one week of harvest on Wheat, Rye and Oats; within 2 weeks of harvest on Barley. Do not pasture dairy animal or meat animals on treated fields.

CAUTION

Harmful if swallowed. Avoid prolonged breathing of dust mist. Avoid contact with eyes, skin, and clothing. Wash thoroughly. Avoid contamination of food or foodstuffs. Keep away from children and domestic animals. Do not use on cucurbits. Do not use on feed or forage.

This product will kill fish and wildlife. Keep out of any body of water.

To prevent re-use completely empty bags and dispose of waste pesticide by burying. Burn bags immediately and stay out of smoke. Bury ashes.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.