



148-1025

CAUTION

HARMFUL IF INHALED OR SWALLOWED. Avoid Prolonged or Repeated Breathing of Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Unnecessary Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD. FEED. FEEDING TROUGHES AND WATERING RECEPTACLES. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application.

NON-WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use. *Trade-mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

3%

SEVIN* DUST

ACTIVE INGREDIENTS:	BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)	3.0%
INERT INGREDIENTS	97.0%
	<u>100.0%</u>

CAUTION KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-104

P-C
PLANTERS CHEMICAL COMPANY
Division of Thompson-Hayward Chemical Company
NORFOLK, VA. FAYETTEVILLE, N. C. ALBANY, GA.

DIRECTIONS FOR USE

POTATOES: To control Colorado Potato Beetle—Dust at the rate of 15 to 20 lbs. to the acre. To control Leafhopper and Flea Beetle — Dust at the rate of 35 lbs. to the acre. Apply dust thoroughly to upper and lower leaf surfaces. Repeat applications, if necessary, at 7 - 10 day intervals.

BEANS: To control Mexican Bean Beetle and Leafhopper — Dust at the rate of 15 to 20 lbs. to the acre. Apply dust thoroughly to upper and lower leaf surfaces. Repeat applications, if necessary, every 7 - 10 days. Can be applied the day of harvest. Will not control insects if used with lime or other highly alkaline materials. Do not graze treated fields or feed trash to dairy or meat animals.

TOMATOES: To control Tomato Fruitworm and Flea Beetle — Dust at the rate of 30 to 50 lbs. per acre. Repeat applications, if necessary, at 7 - 10 day intervals. Can be applied the day of harvest.

CUCUMBERS AND SUMMER SQUASH: To control Striped and Spotted Cucumber Squash Bug, Flea Beetle, Pickleworm and Melonworm — Dust at the rate of 30 to 35 lbs. per acre. Can be applied the day of harvest.

PEPPERS AND EGGPLANT: To control Flea Beetle, Colorado Potato Beetle and Fall Armyworm — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, at 7 - 10 day intervals. Can be applied the day of harvest.