

PLANTERS 7½% CAPTAN DUST

ACTIVE INGREDIENT	
Captan*	7.50%
INERT INGREDIENTS	92.50%
	100.00%

* 1-(2-chlorophenyl)-3-(4-cyclohexene-1,2-dicarboximide)

DIRECTIONS:

STRAWBERRIES: To control Gray Mold or Botrytis Blight — Apply at the rate of 35 to 40 lbs per acre at one-week intervals from the time plants start to grow actively in the spring.

POTATOES (SEED PIECES): To control Seed Rot, Damping off, Verticillium Wilt and Brown Eye Disease — Use at the rate of 10 oz. per bushel cut seed pieces; (1 lb. per 100 lbs. cut seed pieces). Apply in convenient container or by dust attachment over belt. Cut pieces should be treated within 6 hrs. of cutting. If planting is to be delayed more than 1 or 2 days, the treated pieces should be stored for 2 to 3 days in open crates before bagging.

Store in cool, dry place. Protect from excessive heat.

Burn bag immediately, when empty. Stay out of smoke.

CAUTION

Avoid inhaling dust. Wear a respirator while handling. Avoid prolonged or repeated contact with skin. Avoid contamination of feed and foodstuffs.

Do not use on fish. Keep out of any body of water. Do not apply when weather conditions favor drift or dust from areas treated. Containers should be disposed of by punching holes in them and burying with wastes, or by burning. Keep out of smoke!


Seller makes no warranty of any kind, express or implied, concerning the use of this product.

Buyer assumes all risk of use or handling whether in accordance with directions or not.

CAUTION KEEP OUT OF REACH OF CHILDREN

USDA Reg. No. 1868-122

NORFOLK, VA



ALBANY, GA.