

**READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNING AND DIRECTIONS; AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.**

**DIRECTIONS FOR USE**

**ASPARAGUS:** Use 20-40 lbs. per acre for asparagus beetle on seedlings or spear. Use 40-80 lbs. per acre for asparagus beetle and Agave cicada on ferns or brush growth in the post-harvest period. Do not apply more often than once every three days, nor within 1 day of harvest.

**BEANS:** (Green beans, Lima beans, snap beans, cowpeas and black-eyed peas) Use 10 lbs. per acre for Mexican bean beetle. Use 20 lbs. per acre for bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, leafhoppers, velvet bean caterpillar, and spotted bean caterpillar. Use 20 to 30 lbs. per acre for armyworms, corn earworms, stink bugs and tarnished plant bug. Use up to the day of harvest.

**CORN:** Use 20 to 40 lbs. per acre for corn earworm, corn rootworm adults, European corn borer, fall armyworm, flea beetle, Japanese beetle, leaf beetles and leafhoppers. For armyworm, corn rootworm, cutworm and forage feeders, apply to entire plant. Repeat as necessary for insects attacking leaves and ears, apply at 2 to 3 days intervals starting when first insects appear and continuing until silks begin to dry. Three or four applications may be required depending on severity of infestation. Timing and good coverage are important for effective control. Use up to the day of harvest. Application of this product to the tassels region of corn during the polished period will seriously reduce bee population.

**CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER AND KOHLRAB:** Use 10 to 20 lbs. per acre for flea beetles and harlequin bugs. Use 20 to 40 lbs. per acre for armyworm and corn earworm. Use 20 to 40 lbs. per acre for small cabbage beetle. Use 5 to 7 days schedule. Use up to 3 days of harvest.

**HEAD LETTUCE AND CARROTS:** Use 10 to 20 lbs. per acre for flea beetle, harlequin bug and leafhoppers. To control armyworm, cutworms use 20 to 30 lbs. per acre in a 5 to 7 day schedule. Use 20 to 40 lbs. per acre for armyworms, corn rootworm, flea beetle, tarnished plant bug, and flea beetle. For control of corn earworm after lettuce heads begin to form, use 40 lbs. per acre in a 5 to 7 day schedule. Use 20 to 40 lbs. per acre for small cabbage beetle. Use 5 to 7 day schedule. Do not use within 3 days of harvest for lettuce, use up to day of harvest on carrots.

**CUCUMBER, MELONS, PUMPKIN AND SQUASH:** Use 10 to 20 lbs. per acre for flea beetle and leafhoppers. Use 20 to 40 lbs. per acre for armyworm, flea beetle, leafhoppers and Japanese beetle. Some leafhoppers may be difficult to control with this product when tender foliage is wet. The presence of unopened flowers may cause some leafhoppers to be difficult to control. Do not use within 3 days of harvest. Use up to the day of harvest.

**POTATO, TOMATO, EGGPLANT AND PEPPERS:** Use 10 to 20 lbs. per acre for flea beetle, flea beetle, flea beetle and leafhoppers. Use 20 to 40 lbs. per acre for European corn

**ORZA:** Use 20 to 40 lbs. per acre for corn earworm and stink bugs. Repeat at 5 to 7 days intervals or as necessary. Use up to day of harvest.

**BLUEBERRIES:** Use 30 to 40 lbs. per acre for blueberry maggot, cherry and cranberry fruitworm, and Japanese beetle. Apply 3 weeks before harvest and repeat 10 days later or as necessary. Use up to the day of harvest.

**STRAWBERRIES:** Use 20 to 40 lbs. per acre for meadow spittlebug, meadow leaf folder and strawberry weevil. Do not apply within 1 day of harvest.

**ALFALFA, CLOVERS, COWPEAS, PEANUTS AND SOYBEANS:** Use 10 to 20 lbs. per acre for beetles and Mexican bean beetle. Use 20 lbs. per acre for alfalfa caterpillar, leaf beetles, cucumber beetle, green cloverworm, Japanese beetle, leafhopper, three cornered alfalfa hopper, thrips, and alfalfa caterpillar. Use 20 to 30 lbs. per acre for armyworm, corn earworm, stink bugs and webworms. Some leaf injury may occur if applications are made when tender foliage is wet. Use up to day of harvest.

**FORAGE, GRASSES AND PASTURE:** Use 20 to 30 lbs. per acre for armyworms, stink bugs and thrips. Use up to the day of harvest.

**SORGHUMS (Including MILO AND GRAIN SORGHUM):** Use 20 to 40 lbs. per acre for armyworm, corn earworms, stink bug and webworms. For optimum insect control on grain sorghum, apply to entire forming heads. Use 30 lbs. per acre for armyworms. Treatment for sorghum midge, corn rootworm and cutworm 3 to 4 days after heads have emerged from soil. Do not use more than 20 days of harvest.

**TOBACCO - In Plant Beds:** For Tobacco Flea Beetle, apply 1 to 2 lbs. per acre in square yard when plants appear. Repeat at 5 to 7 day intervals and just before transplanting.

**TOBACCO - In Fields:** For insect applications for hornworms, apply 15 to 20 pounds per acre in a 5 to 7 day schedule. For flea beetle, apply 20 to 40 lbs. per acre in a 5 to 7 day schedule. Repeat as necessary when tender foliage is wet. Do not use within 3 days of harvest.

**NOTE:** If insect applications are necessary, treat immediately after planting and before pinning. Treated fields may be grazed immediately after it has settled.

Some insect injury may occur on tender forage in the post-harvest period in the absence of several days duration following applications.

**NOTICE — Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this**



# 5% SEVIN\* DUST

<b>ACTIVE INGREDIENTS:</b>	By Weight
Carbaryl (1-naphthyl N-methylcarbamate)	5.0%
<b>INERT INGREDIENTS:</b>	95.0%
	<hr/> 100.0%

## CAUTION KEEP OUT OF REACH OF CHILDREN

**CAUTION: HARMFUL IF INHALED OR SWALLOWED.** Avoid Breathing of Dust. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Contact. Wash Hands and Face Before Eating. Take Shower or Bath After Work. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

**NOTE FOR PHYSICIAN:** Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

\*Trade Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate U. S. Pat. No. 2,903,478