## DIRECTIONS

For the control of leaf spot on peanuts apply at 7 to 10 day intervals beginning when spots first appear on basal leaves at the rate of 15 to 20 pounds per acre.

Armyworms, Velvet Bean Caterpillar and peanut worms on peanuts: begin applications at the rate of 20 to 30 pounds per acre when small worms first appear. During periods of severe infestations rates should be increased.

Thrips on peanuts: begin applications at the rate of 20 to 30 pounds per acre when small thrips first appear. Repeat the application in 7 to 10 days. Apply additional dustings during the season as needed.

5% SEVIN*
85% SULPHUR
PEANUT DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)	5.00%
Sulphur	85.00%
INERT INGREDIENTS:	10.00%
TOTAL	100.00%

*Trade-Mark of Union Carbide Corp. for 1-naphthyl N-methylcarbamate

CAUTION KEEP OUT OF REACH OF CHILDREN.

USDA Reg. No. 1868-15

15%

P-C

PLANTERS CHEMICAL COMPANY
Division of Thompson-Hayward Chemical Company
NORFOLK, VA.   FAYETTEVILLE, N. C.   ALBANY, GA.

CAUTION

Avoid excessive inhalation. Harmful if taken internally. Keep away from children, pets, livestock, feed, and foodstuffs. Do not apply to pastures, forage crops or other edible portions of plants which will be consumed by dairy animals or to animals being finished for slaughter.

This product is toxic to fish and wildlife. Do not apply when weather conditions favor drift from areas treated. Do not contaminate streams, lakes or ponds with this material.

NON-WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.