

DIRECTIONS FOR USE

POTATOES: To control Aphid, Colorado Potato Beetle, Leafhopper, Flea Beetle, Early and Late Blight — Dust at the rate of 35 to 40 lbs. per acre, depending on the size of plants and severity of insect and disease conditions. Apply thoroughly to upper and lower leaf surfaces. Repeat applications at 7-10 day intervals. Do not apply within 5 days of harvest.

TOMATOES: To control Aphid, Tomato Fruitworm, Flea Beetle, and Early and Late Blight — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, within 10 days of harvest.

CUCUMBERS AND SUMMER SQUASH: To control Aphid, Stained and Spotted Cucumber Beetle, Squash Bug, Flea Beetle, Pickleworm, Melonworm, Downy Mildew, Anthracnose, Bacterial Wilt and Alternaria Leaf Spot — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, at 7-10 day intervals. Do not apply within 15 days of harvest.


PEPPERS AND EGGPLANT: To control Aphid, Flea Beetle, Colorado Potato Beetle, Fall Armyworm, Leaf Spot and Blight — Dust at the rate of 30 to 35 lbs. per acre. Repeat applications, if necessary, at 7-10 day intervals. Do not apply within 15 days of harvest.

DANGER

May be fatal if swallowed, inhaled, or absorbed through skin. Avoid breathing dust. Rapidly absorbed through skin. Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Workers entering fields within 24 hours after application should wear protective clothing. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not re-use empty containers. Destroy empty bags and waste pesticide by burning. Stay out of smoke and fumes. Bury ashes in non-crop lands away from water supply.



ACTIVE INGREDIENTS.		BY WEIGHT
Carbaryl (1-naphthyl N-methylcarbamate)		3.0%
Parathion**		1.0%
Copper Oxide		8.45%
(Copper expressed as Metallic		7.0%)
INERT INGREDIENTS:		87.55%
TOTAL		100.0%

DANGER KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-194

NOT FOR USE OR STORAGE IN OR AROUND THE HOUSE



**THOMPSON-HAYWARD
CHEMICAL COMPANY**

PLANTERS CHEMICAL DIVISION

NORFOLK, VA.

FAYETTEVILLE, N. C.

ALBANY, GA.

FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING! If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets each (1/100 gr.) at once. Remove patient immediately from the area where Parathion is present. Remove all contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of parathion. If swallowed, induce vomiting by giving warm salty or soapy water.

ANTIDOTE

ATROPINE AND 2-PAM ARE THE EMERGENCY ANTIDOTES FOR METHYL PARATHION POISONING. Keep on hand an emergency supply of 25 Atropine tablets, grains 1/100. In case of poisoning take 3 tablets (grs. 1/100) and call physician at once.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

PHYSICIANS' NOTE

Parathion is a Cholinesterase inhibitor and treatment of severe poisoning should include the use of atropine.

Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT

Give atropine, preferably by injection, grains 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

*Trade-Mark of Union Carbide Corporation for the active ingredient Carbaryl (1-naphthyl N-methylcarbamate).

**O,O-diethyl O-p-nitrophenyl thiophosphate.