

## DIRECTIONS FOR USE ON SHADE TOBACCO

**IN THE BED:** For the control of Blue Mold, Aphids, Thrips, Flea Beetles, Vegetable Weevils and Slugs. Apply at the following rates. Apply 2 to 4 lbs. per 100 square yards when plants are the size of a dime and before Blue Mold appears. Dust every 3 days.

**IN THE SHADE:** For the control of Aphids, Thrips, Flea Beetles, Budworms, Hornworms and Cabbage Loopers. Apply weekly, or as needed, at 20 to 30 lbs. per acre, depending on size of plants.

"Florida Shade Dust" is formulated with "Lignoflex" a special Pecan Hull Flour Diluent approved for use on Shade Tobacco.

### CAUTION

Do not apply to tobacco within 5 days of priming or 15 days of cutting. Workers should wear gloves where applications have been made within 30 days of cutting. Avoid plant juices coming in contact with the skin or other parts of the body when engaged in cutting the crop.

### WARNING


May be fatal if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin! Do not breathe dust. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Workers entering a treated field within 5 days of an application should wear protective clothing. Do not contaminate feed or foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. Do not apply when weather conditions favor drift from areas treated.

This product is highly toxic to bees exposed to direct treatment or residues on crop. Protective information may be obtained from Cooperative Agricultural Extension Service.

Do not graze treated fields.

To prevent re-use completely empty bags and dispose of waste pesticide and bags by burying.



<b>ACTIVE INGREDIENTS:</b>	
Parathion (O,O-diethyl O-p-nitrophenyl thiophosphate)	2.00%
Endosulfan (Hexachlorohexahydromethano-2,1,3-Denzodioxathiepin oxide)*	3.00%
Zineb (Zinc Ethylene Bisdithiocarbamate)	13.00%
<b>INERT INGREDIENTS:</b>	82.00%
<b>TOTAL</b>	<b>100.00%</b>
*Thiodan - U. S. Patent No. 2,799,685	

**DANGER KEEP OUT OF REACH OF CHILDREN**

USDA Reg. No. 1868-240

## FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING. If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets each (1-100 gr.) at once. Remove patient immediately from the area where Parathion is present. Remove all contaminated clothing and wash clean with plenty of soap and water to remove all traces of Parathion. If swallowed, induce vomiting by giving warm salty or soapy water.

## PHYSICIANS' NOTE

Parathion is a cholinesterase inhibitor and treatment of severe poisoning should include the use of atropine; Endosulfan produce (s) central nervous system excitation which responds to the barbiturate acid derivatives.

Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

## TREATMENT

Give atropine, preferably by injection, grains 1-100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

## ANTIDOTE

ATROPINE IS AN ANTIDOTE. Consult Physician For Emergency Supply.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.