

PLANTERS

**DIRECTIONS FOR USE**

Apply 2 1/2 lbs. of this material per acre unless otherwise noted to control the following insects on the crops shown below. Apply when the air is calm by ground machinery. Thorough coverage of the plant is necessary for effective control. Observe days interval between last application and harvest indicated by number in parenthesis following crop.

**TOMATOES (10):** For the control of Aphids, Colorado Potato Beetle, Flea Beetle and Whitefly.

**HEAD LETTUCE (14):** For the control of Cabbage Looper, Green Peach Aphid, Diamond-Back Moth, Larvae, and Imported Cabbage Worm. Note: Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed crop refuse to livestock.

**MELONS (14):** For the control of Aphids and Cucumber Beetles. **PEPPERS (15):** For the control of Flea Beetle and Green Peach Aphid.

**POTATOES (10):** For the control of Flea Beetle, Colorado Potato Beetle, Three Lined Beetle, Leafhopper, Aphids and Plant Bugs, Potato Tuberworm, and Whitefly.

**CAULIFLOWER:** For the control of Cabbage Looper, Imported Cabbage-worm, Diamond-back Moth Larvae, and Cross-striped Cabbage-worm. Note: Do not apply after edible parts begin to form.

**BROCCOLI (7), CABBAGE (10):** For the control of Cabbage Aphids, Cabbage Looper, Imported Cabbage-worm, Diamond-back Moth Larvae, Cross-striped Cabbage-worm, Flea Beetle, Harlequin Bug, and Stink Bug.

**SUCCULENT BEANS:** Mexican Bean Beetle, Leafhopper, Aphid, Leaf Roller, Armyworms, Leaf Miner - 15 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated thrashings to livestock.

**SOYBEANS:** Green Stink Bug, Mexican Bean Beetle, Velvet Bean Caterpillar - 15 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated thrashings to livestock.

**SUGAR BEETS (17):** Green Peach Aphid, Sugar Beet Web-worm - 33 lbs. per acre. Do not feed treated sugar beet tops to livestock.

**MELONS (14), CUCUMBER (15), SQUASH (15), PUMPKINS (15):** Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer, Stink Bugs, Melonworms, Pickleworm, Serpentine Leaf Miner. Do not apply to cucurbits unless plants are dry. Make no application before plants start to vine.

**TOBACCO (15):** Flea Beetles, Hornworm, Budworm, Green Peach Aphid. Thorough coverage is essential for good control. Make no application within 7 days of priming. Avoid getting wet with rain, dew, or water during or priming.

ACCEPTED

198-1076  
FERTILIZERS & CHEMICALS  
DIVISION OF THE  
FARMERS' COOPERATIVE UNION  
OF AMERICA  
ST. LOUIS, MO.  
DISTRIBUTED BY  
CUMMINGS

**PARATHION 2  
THIODAN® 3  
DUST**  
INSECTICIDE

®Thiodan is a registered Trademark of Farbwerke Hoechst A. G.

**POISON**



<b>ACTIVE INGREDIENTS:</b>	
*Parathion	2.0%
*Endosulfan (Hexachloro-hexahydro-methano-2,4,3-benzodioxathiepin oxide)	3.0%
<b>INERT INGREDIENTS:</b>	
	95.0%
<b>TOTAL</b>	<b>100.0%</b>
*O,O-Diethyl O-p-nitrophenyl thiophosphate	
*Thiodan, U. S. Patent No. 2,799,685	

**DANGER KEEP OUT OF REACH OF CHILDREN**

USDA REG. No. 1868-

T-H

**DANGER - POISON**

**Antidotes:** If swallowed, give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Have victim lie down and keep quiet. Call a Physician Immediately! If on skin, in case of contact, remove contaminated clothing and immediately wash skin with soap and water. Call a Physician Immediately!

**ATROPINE IS ANTIDOTAL**

Keep on hand an emergency supply of 20 tablets (grams 1.100). In case of accidental poisoning take 3 tablets (grs. 1.100) and call physician at once.

**Physician:** Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

**WARNING**

**WARNING:** May be Fatal if Swallowed, Inhaled or Absorbed Through Skin! Rapidly Absorbed Through Skin! Do not get in eyes, on skin, or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms, and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as specified on this label.

**DISPOSAL WARNING:** Do not reuse the container. Completely empty the contents and bury the unused chemical with empty bags at least 12 inches deep in an isolated location away from water supplies.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Dealers Should Sell in Original Packages Only.

USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH THE DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES OR OTHER UNDESIRABLE RESULTS.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACCEPTED WITH COMMENTS