

DIRECTIONS

This material is intended for use as a dust on Peanuts for the control of Cercospora Leafspot, Aphids, and Red Spider Mites. Application should be made at the rate of 25 to 35 lbs. per acre with a dusting machine capable of depositing a thin uniform coating of dust on all above ground plant surfaces. Begin applications when disease first appears in the area and repeat at 14 day intervals for a total of three to four applications.

CAUTION

Applications of dusting or spraying materials containing sulphur should not be made just prior to or during periods of excessively high temperatures. USE OF THIS MATERIAL IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH DIRECTIONS MAY PRODUCE PLANT INJURY, EXCESSIVE RESIDUES, OR OTHER UNDESIRABLE RESULTS.

Not for use or storage in or around the home.

DANGER

Poisonous if swallowed, inhaled or absorbed through skin. Rapidly absorbed through skin! Do not breathe dust. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Workers entering a treated field within 5 days of an application should wear protective clothing. Do not contaminate feed or foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use.

Toxic to wildlife. Do not contaminate any body of water with this material by direct application, cleaning of equipment or disposal of wastes and containers; nor apply to area not specified on label. Birds feeding on treated areas may be killed.

Do not graze dairy animals or animals being finished for slaughter on fields which have received late application of this insecticide.

To prevent re-use completely empty bags and dispose of waste pesticide by burying. Burn bags immediately and stay out of smoke. Bury ashes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.



148-1098

ACCEPTED

Sulfur - Copper
2 1/2% M.P. Dust

POISON



ACTIVE INGREDIENTS:

| | |
|--|---------|
| Sulfur | 70.00% |
| Copper (in copper oxide form) expressed as metallic | 3.40% |
| *O,O-dimethyl O-p-nitrophenyl phosphorothioate | 2.50% |
| INERT INGREDIENTS: | 24.10% |
| TOTAL | 100.00% |

*Otherwise known as Methyl Parathion

DANGER KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1869-251

SEE ANTIDOTE STATEMENT AND OTHER WARNINGS ON SIDE PANEL



**THOMPSON-HAYWARD
CHEMICAL COMPANY**

PLANTERS CHEMICAL DIVISION

NORFOLK, VA.

FAYETTEVILLE, N. C.

ALBANY, GA.

FIRST AID TREATMENT

CALL A PHYSICIAN AT ONCE IN ALL CASES OF SUSPECTED PARATHION POISONING! If symptoms or signs of poisoning include blurred vision, abdominal cramps and tightness in the chest, don't wait for a doctor but give two atropine tablets each (1/100 gr.) at once. Remove patient immediately from the area where Parathion is present. Remove all contaminated clothing and wash the skin clean with plenty of soap and water to remove all traces of parathion. If swallowed, induce vomiting by giving warm salty or soapy water.

ANTIDOTE

ATROPINE AND 2-PAM ARE THE EMERGENCY ANTIDOTES FOR METHYL PARATHION POISONING. Keep on hand an emergency supply of 25 Atropine tablets, grains 1/100. In case of poisoning take 3 tablets (grs. 1/100) and call physician at once.

NEVER GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON.

PHYSICIANS' NOTE

Parathion is a Cholinesterase inhibitor and treatment of severe poisoning should include the use of atropine.

Warning symptoms include weakness, headache, tightness in the chest, blurred vision, non-reactive pin-point pupils, salivation, sweating, nausea, vomiting, diarrhea and abdominal cramps.

TREATMENT

Give atropine, preferably by injection, grains 1/100 two or three tablets at once and parenterally or orally every hour as required up to 30 tablets or until pupils dilate. Never give morphine. Clear chest by postural drainage. Artificial respiration or oxygen administration may be necessary. Observe patient continuously for 48 hours. Repeated exposure to cholinesterase inhibitors may, without warning, cause prolonged susceptibility to very small doses of any cholinesterase inhibitor. Allow no further exposure until time for cholinesterase regeneration has been allowed as determined by blood tests.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.