

DIRECTIONS FOR USE

Apply when insects first appear and repeat as required to maintain effective control. Coverage of both upper and lower leaf surfaces is essential for good control. Observe all precautions. If insect control is required beyond the following use patterns, consult the control program with other suitable practices. Observe days interval between last application and harvest indicated by number in () following crop.

APPLES (30): Apple Aphid - 50 lbs. per acre. Unlimited number of applications.

OR

(21): Apple Aphid - 50 lbs. per acre. Do not make more than 3 applications during the fruiting period.

Pomace from treated apples should not be fed to livestock.

APRICOTS (30), NECTARINES (30), PEACHES (30): Green Peach Aphid - 50 lbs. per acre. Do not make more than 2 applications during the fruiting period.

SUCCULENT BEANS: Mexican Bean Beetle - 18 to 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock. Do not apply more than three times per season. Do not allow livestock to graze in treated fields. Do not use on lima beans that are to be removed from the field for processing.

CAULIFLOWER: Imported Cabbage Worm, Cabbage Looper, Diamondback Moth Larvae, Cross-striped Cabbageworm, Flea Beetles, Harlequin Bugs, Stink Bugs - 25 to 33 lbs. per acre. Do not apply after edible parts begin to form. Close trim other leaves prior to marketing.

BROCCOLI (7), CABBAGE (7): Imported Cabbage Worm, Cabbage Looper, Diamondback Moth Larvae, Cross-striped Cabbageworm, Flea Beetles, Harlequin Bugs, Stink Bugs - 25 to 33 lbs. per acre.

CUCUMBERS (0): Aphids, Cucumber Beetles, Striped Cabbage Flea Beetle - 25 to 33 lbs. per acre.

EGG PLANTS (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Green Stink Bug, Whitefly - 17 lbs. per acre.

LETTUCE, HEAD (14): Cabbage Looper, Diamondback Moth Larvae, Green Peach Aphid, Imported Cabbage Worm - 25 to 33 lbs. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves at harvest. Do not feed refuse to livestock.

MELONS (0): Aphids, Cucumber Beetle, Striped Cabbage Flea Beetle - 25 to 33 lbs. per acre; Cabbage Looper, Omniverous Leaf Roller - 33 lbs. per acre.

ORNAMENTALS: Aphids, Whitefly - Dust lightly, but uniformly. On Chrysanthemums best results will be obtained if applied before plants flower. Do not apply to Birch Trees.

PEPPERS (1): Flea Beetle, Green Peach Aphid - 33 lbs. per acre. Pepper Maggot - 17 lbs. per acre.

POTATOES (0): Aphids, Colorado Potato Beetle, Plant Bugs (including Shield Shaped Bugs), Potato

(CONTINUED ON RIGHT PANEL)

PLANTERS

REGISTERED
THIODAN[®] 3
DUST
INSECTICIDE

ACTIVE INGREDIENT:

*Endosulfan (Hexachlorohexahydromethano-2,4,3-benzodioxathiepin oxide) 3.0%

INERT INGREDIENTS: 97.0%

TOTAL 100.0%

*Thiodan U. S. Pat. No. 2,799,685.

*Thiodan is a registered trademark of Farbwerke Hoechst A. G.

CAUTION KEEP OUT OF REACH OF CHILDREN

USDA REG. No. 1868-253



Flea Beetle, Potato Leafhopper, Potato Tubeworm, Southern Armyworm - 30 to 35 lbs. per acre; European Corn Borer, Potato Psyllid - 35 lbs. per acre; False Chinch Bug - 35 lbs. per acre; Three-lined Potato Beetle - 17 lbs. per acre.

PUMPKINS (0): Aphids, Cucumber Beetles - 25 to 33 lbs. per acre.

SAF FLOWER: Green Peach Aphid - 35 lbs. per acre. Do not apply after flower heads open.

SOYBEANS: Green Stink Bug, Mexican Bean beetle - 20 lbs. per acre. Do not apply after pods begin to form. Do not feed treated threshings to livestock.

SQUASH (0): Aphid, Cucumber Beetles, Striped Cabbage Flea Beetle, Squash Vine Borer - 25 to 33 lbs. per acre. For Vine Borer control, apply weekly to flower buds, stems and vines beginning when moths first appear.

STRAWBERRY (4): Meadow Spittlebug, Strawberry Aphid - 33 lbs. per acre. Do not apply at intervals of less than 15 days or more than twice during any 35 day period when fruit is present.

SUGAR BEETS (0): Green Peach Aphid, Sugar Beet Webworm - 33 lbs. per acre. Do not feed treated sugar beet tops to livestock.

TOBACCO (5): Flea Beetles, Hornworm, Budworm, Green Peach Aphid - 25 to 30 lbs. per acre. Thorough coverage is essential for good control. **SEED BED**: Use 2 lbs. per 100 sq. yds.

TOMATOES (FIELD AND GREENHOUSE) (1): Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Peach Aphid, Hornworms, Whitefly - 17 lbs. per acre. Tomato Fruitworm, Yellow-striped Armyworm - 35 lbs. per acre. Stink Bugs - 20 to 33 lbs. per acre. Cabbage Looper - 25 lbs. per acre.

CAUTION

Harmful if swallowed, inhaled or absorbed through skin. Avoid breathing of dust. Avoid contact with skin, eyes or clothing. In case of contact with skin or eyes, flush with plenty of water; for eyes, get medical attention. Wash with soap and water after handling and before eating or smoking; wear clean clothing. Avoid contamination of feed and food-stuffs. Do not apply or allow to drift to areas occupied by unprotected humans or beneficial animals.

This product is toxic to fish and wildlife. Do not contaminate any body of water nor apply when conditions favor drift from the crop area treated. Apply this product only as specified on this label. This product is toxic to bees and should not be applied when bees are actively visiting the area.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer as-