

number in parenthesis following crop.

FOR AIRPLANE APPLICATIONS: Use approximately 3 to 10 gallons of water per acre consistent with crop growth and good coverage.

BLOCCOLI (7), CABBAGE (10): To control Cabbage, Aphids, Cabbage Looper, Imported Cabbageworm, Diamond-back Moth, Larvae, Cross-striped Cabbageworm, Flea Beetle, Harlequin Bug, Stink Bug use 1½ to 2 qts. per acre.

CAULIFLOWER: To control Cabbage Looper, Imported Cabbageworm, Diamond-back Moth, Larvae, Cross-striped Cabbageworm use 1 to 1½ qts. per acre. To control Flea Beetle, Harlequin Bug, Stink Bug use 1½ qts. per acre. Do not apply after edible parts begin to form.

CUCUMBERS (15), PUMPKINS (15), SUMMER SQUASH (15), WINTER SQUASH (15): To control Aphids, Cucumber Beetles use 1 to 2 qts. per acre. Do not apply before vining.

HEAD LETTUCE (14): To control Cabbage Looper, Green Peach Aphid, Diamond-back Moth Larvae, Imported Cabbage Worm use 1½ to 2 qts. per acre. Do not make more than 3 applications after thinning. Remove wrapper leaves. Do not feed crop refuse to livestock.

MELONS (7): To control Aphids, Cucumber Beetles use 1 to 2 qts. per acre. To control Cabbage Looper, Omniverous Leaf Roller use 2 qts. per acre.

PEPPERS (15): To control Flea Beetle, Green Peach Aphid use 1 qt. per acre.

POTATOES (5): To control Flea Beetle, Colorado Potato Beetle, Leafhopper, Aphids, Southern Armyworm, Green Stink Bug, Leaf-footed Plant Bugs and other Plant Bugs, Potato Tuberworm, Whitefly use 1 to 2 qts. per acre. To control False Chinch Bug use 2 qts. per acre. To control Potato Psyllid use 1½ to 2 qts. per acre.

SUGAR BEETS (15): Green Peach Aphid, Sugar Beet Webworm, Meadow Spittlebug, Leafhoppers, Blister Beetles, Flea Beetles, Leaf Miner - Use 2 quarts per acre. Apply when insects first appear and repeat as required. Do not feed treated beet tops to livestock.

TOMATOES (10): To control Aphids, Blister Beetle, Colorado Potato Beetle, Flea Beetle, Green Stink Bug, Whitefly use 1 qt. per acre. To control Hornworm use 1 to 2 qts. per acre.

COTTON: To control Spider Mites, (Desert Spider Mite, Tumid Spider Mite, Two-spotted Mite), Boll Weevil, Aphids, Cotton Leafworm, Lygus Bug, Bollworm, Cabbage Looper, use 1½ qts. per acre. **NOTE:** Do not apply after 25% of the bolls open. Do not graze dairy or meat animals in treated fields.

TOBACCO: To control Budworm and Tobacco Suckfly, use 2 pints per acre. To control Cabbage Looper, Flea Beetle, Green Peach Aphid and Hornworm, use 1 to 2 quarts per acre. **NOTE:** Do not apply within 5 days of priming or within 15 days of cutting. Do not enter fields within 24 hours of application. Protective clothing should be worn during applications.

TOBACCO (PLANT BED TREATMENT): Green June Bug Larvae - 1 pint per 100 gallons. At first sign of larvae activity, apply as a drench using 1 gallon of spray mixture per square yard of plant

PARATHION 1 THIODAN 2 Emulsifiable Concentrate INSECTICIDE

ACTIVE INGREDIENTS:

Parathion	11.5%
Endosulfan (Hexa-hydrohexahydro-methano-2,4,3-benzodioxathiepin oxide)	22.8%
Xylene base aromatic petroleum solvent	60.0%
INERT INGREDIENTS:	5.7%

O, O-Diethyl O-p-trophenyl phosphosphate Thiodan, U. S. Patent No. 2,799,655	100.0%
---	--------

Contains 1.0 lb. of actual Parathion and 2.0 lbs. of actual Thiodan per gallon.

*Thiodan is a registered trademark of Farbwerke Hoechst A. G.

DANGER KEEP OUT OF REACH OF CHILDREN

USDA Reg. No. 1868-255

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of such use.

See antidote statement and other warnings on the side panel.



water.

ATROPINE AND 2-PAM ARE THE EMERGENCY ANTIDOTES FOR METHYL PARATHION POISONING. Keep on hand an emergency supply of 25 Atropine tablets, grains 1/100. In case of poisoning take 3 tablets (grs. 1/100) and call physician at once.

Physician: Parathion is a cholinesterase inhibitor and treatment of poisoning should include the use of atropine; Thiodan product(s) central nervous system excitation which responds to the barbituric acid derivatives. Enough atropine must be given in the shortest practical time to produce dilation of pupils. Patient must be under constant supervision and additional atropine may be necessary without delay. Clear chest by postural drainage. Use artificial respiration and oxygen if patient becomes comatose. Use of airway may be necessary.

DANGER

Poisonous if swallowed, inhaled or absorbed through skin! Rapidly absorbed through skin. Do not get in eyes, on skin or on clothing. Wear natural rubber gloves, protective clothing and goggles. In case of contact, wash immediately with soap and water. Wear a mask or respirator of a type passed by the U. S. Department of Agriculture for Parathion protection. Keep all persons and beneficial animals out of operating area or vicinity where there may be danger of drift. Vacated areas should not be re-entered until drifting insecticide and volatile residues have dissipated. Do not contaminate feed and foodstuffs. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before re-use. Workers entering treated fields within 24 hours should wear protective clothing.

This product is toxic to fish and wildlife. Keep out of any body of water. Apply this product only as birds feeding in treated areas may be killed. Do not apply when weather conditions favor drift from a gas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DISPOSAL WARNING: Do not reuse the container. Completely empty the contents and bury the unused chemical and empty container at least 18 inches deep in an isolated location away from water supplies.

WARNING - FLAMMABLE

Keep Away From Heat and Open Flame. Keep Container Closed.

Leaking Packages Should Be Removed To A Safe Place.

Destroy Containers When Empty.

CAUTION: Do Not Use In Undiluted Form. Do not Store At Temperatures Below 20° F.

Dealers Should Sell in Original Packages Only.

Do not allow this material to drift onto neighboring crop, or non-crop areas, or use in a manner or at a time other than in accordance with directions, because plant injury, excessive residues, or other