DED-WEED LFB

MIXING INSTRUCTIONS

Determine the amount of Ded-Weed LFB to be used in the sprayer by multiplying the number of acres to be treated by the quantity of Ded-Weed LFB recommended per acre. Add this amount of Ded-Weed LFB to one-half the water required for coverage. Mix well, then add remaining water. Continue mixing until spray mixture is uniform. Best results are obtained by treatment when weeds are growing vigorously.

FOR PERENNIAL WEEDS IN PASTURE, RANGELANDS AND NON-CROP LANDS SUCH AS FENCE ROWS, DRAINAGE DITCH BANKS, RIGHT-OF-WAYS

WEEDS	Amount Ded-Weed LFB Per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
CANADA THISTLE AND PERENNIAL SOW THISTLE	1½ pints	3/4 lb.	Apply during early bud stage and again when rosettes are formed.	Use 1½ pints per acre, each spraying.
BINDWEED	1½ pints	3/4 lb	Apply bud to bloom stage.	Re-apply in late fall
E TOP RY SS	1 to 1½ pints	½ to 3/4 lb	In spring when growing actively and foliage fully expanded.	Re-treat fall resettles with 1 to 2 quarts per acre
RUSSIAN KNAPWEED	1½ pints	3/4 lb	In spring before bloom	Stops top growth and prevents seed production. Re-treat as necessary

Hard to-kill weeds such as Canadian Thistle may not be controlled by one application but top growth and re-seeding usually will be prevented. Repeated treatments generally give good control.

Ded-Weed LFB may be mixed with liquid nitrogen to accomplish both a weeding and feeding of grain crops in one operation. See mixing directions below.

Ded-Weed LFB - Liquid Nitrogen Combination Sprays. Fill the spray tank half full with the liquid nitrogen. Start the agitator. Add Ded-Weed LFB in full quantity to cover the number of acres the full tank will cover. Follow warnings and recommendations for amount Ded-Weed LFB to use. Continue agitation of the spray solution. Complete filling of tank with fertilizer. Begin the spraying and fertilizing application. It is necessary to continue good agitation from the time Ded-Weed LFB is added until the tank is empty. Mix only one tank at a time. Apply immediately. Note: Do not spray during or immediately following cold weather.

control of resistant species of brush and certain timberland applications use other Thompson-Hayward products such as 2,4-D LV-6 Brush Kil and Ded-Weed LV-6 Brush Kil.

Do not reuse empty drum. Return to drum reconditioner or destroy by perforating or crushing and burying in a safe place away from water supplies.

NOTICE: ***Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.***

CAUTION **Keep Out of Reach of Children**

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS

2,4-D 4 LBS. PER GALLON

FORMULATED FOR USE IN LIQUID NITROGEN LOW VOLATILE ESTER

ACTIVE INGREDIENTS:	
2,4-dichlorophenoxyacetic acid isooctyl ester	67.0%
(Equivalent to 2,4-dichlorophenoxyacetic acid 44.4%)	
INERT INGREDIENTS	33.0%
	100.0%

EPA Reg. No. 148-1117

CG-1-73-A

THOMPSON HAYWARD CHEMICAL COMPANY

Dec. 3, 1974

148-1117

8 9/3

Pesticides Registration Division, EPA

DIRECTIONS

BEFORE USE READ DIRECTIONS, CAUTION AND NOTICE STATEMENTS CAREFULLY

WEED CONTROL IN CROPS

BROADLEAF WEEDS IN	Amount Ded-Weed LFB per Acre	Amount 2,4-D Acid Per Acre	TIME OF APPLICATION	REMARKS
SPRING WHEAT RYE, BARLEY	½ to 2/3 pint	1/4 to 1/3 lb	Apply full tiller to boot stage and after milk stage to harvest	Do not treat in fall. Do not apply from boot to milk stage or in seeding stages. See single asterisk* below for additional remarks
OATS	½ pint	1/4 lb.		
FALL WHEAT	½ to 1 pint	1/4 to ½ lb		
FIELD CORN PRE-EMERGENCE	1 to 1½ quarts	1 to 1½ lbs	Any time after planting, preferable 2 to 5 days, but before emergence	Do not apply on sandy soil. Use amounts in 5 to recommended 20 gallons of water per acre
CORN SUNFLOWER POST-EMERGENCE	3/8 to ½ pint	3/16 to 1/4 lb	When weeds are small and crops 4 to 18 inches tall	Use drop nozzles on crop more than 8 inches tall to keep spray off foliage
PASTURES RANGELAND	1½ pint	3/4 lb	When annual or perennial broadleaf weeds are growing actively, fall or spring, perennials near bud stage	Legumes present may be damaged. Do not graze dairy animals on treated areas within 7 days after application. Do not apply to seedling grasses. Do not apply when grass is in early boot to milk stage. When grass is grown for seed production

Do not forage or graze treated grain fields within 2 weeks after treatment with 2,4-D. Do not feed treated straw within [illegible]

FOR DILUTION CALCULATIONS—
1 quart contains 1 lb. 2,4-D acid

PARTIAL LIST OF WEEDS CONTROLLED BY DED-WEED LFB

Carpet Weed, Lambs Quarters, Pigweed, [illegible]

[illegible] temperatures [illegible] be reduced to [illegible] 60°F, before spraying.

CAUTION

[illegible]

are present.