

Apply a 1% Diazinon spray to shelves and cupboards where food containers suspected of being infested are stored. Remove all food and shelving paper before spraying. Do not replace the food packages until spray has dried.

Cover shelves with shelving paper. This treatment is supplemental to good housekeeping, since it will have no effect on insects breeding within food packages. Infested food packages should be destroyed.

FOOD HANDLING ESTABLISHMENTS CRACK AND CREVICE TREATMENT ONLY.—Places other than private residences, in which food is held, processed, prepared and/or served.

For control of Resistant and Non-Resistant Cockroaches (Waterbugs), Ants, Spiders, Silverfish, and as an aid in the control of Pantry Pests—exposed stages Sawtooth Grain Beetle, Flour Beetle, Rice Weevil, Cigarette Beetle, Drugstore Beetle, Indian Meal Moth.

FOOD AREAS—Includes receiving, serving, storage (Dry, Cold, Frozen, Raw), packaging (cleaning, slicing, cooking, grinding), edible waste storage, enclosed processing systems (mills, dairies, edible oils, syrups). Serving areas (when food is exposed and facility is in operation also would be considered a food area).

Apply 1/2%-1% Diazinon spray as a thin stream to crack and crevices only where insects hide or through which they may enter the building. Includes expansion joints, spaces between equipment and floors, and openings leading to voids and hollow spaces in walls, equipment legs and bases, conduits, motor housings, junction or switch boxes. Prior to application de-energize electrical equipment. After application allow time for oil to evaporate before starting motors or placing switches in the "ON" position.

NOTE: Use equipment capable of delivering a pin stream of insecticide.

CARE SHOULD BE TAKEN TO AVOID DEPOSITING THE PRODUCT ONTO EXPOSED SURFACES OR INTRODUCING THE MATERIAL INTO THE AIR. AVOID CONTAMINATION OF FOOD OR FOOD PROCESSING SURFACES.

APPLICATIONS OF THIS PRODUCT IN THE FOOD AREAS OF FOOD HANDLING ESTABLISHMENTS, OTHER THAN AS A CRACK AND CREVICE TREATMENT IS NOT PERMITTED.


DO NOT ALLOW CHILDREN IN TREATED AREAS UNTIL SURFACES ARE DRY.

Brown Dog Ticks in Buildings: Make thorough application of a 1/2% Diazinon spray in infested areas around baseboards, window and door frames, wall cracks, sleeping quarters of household pets, and localized areas of floors and floor covering. Fresh bedding should be placed in animal quarters following treatment. Do not treat animals with this formulation.

WARNING

May be fatal if swallowed. May be absorbed through skin. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Do not reuse empty container. Destroy by perforating or crushing and burying in a safe place.

**DO NOT USE, POUR, SPILL, OR STORE
NEAR HEAT OPEN FLAME.**



DIAZINON® OS-4 INSECTICIDE

FOR USE BY PEST CONTROL OPERATORS ONLY.
FOR DILUTION WITH INSECTICIDE BASE OILS.

WARNING: Keep Out of Reach of Children

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS.

ACTIVE INGREDIENTS:

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl)
phosphorothioate 48.7%
Aromatic petroleum derivative solvent 44.0%

INERT INGREDIENTS: 7.3%
Total: 100.0%


DIAZINON OS-4 contains 4 lbs. Diazinon per gallon.
DIAZINON® is a registered trademark of CIBA-GEIGY CORP.
U.S. Patent No. 2,754,243

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 148-1129

CG-8-74

EPA EST No. 148-1L-1



**THOMPSON-HAYWARD
CHEMICAL COMPANY**

P O BOX 2383 KANSAS CITY KANSAS 66110

DIAZINON OS-4 is for pest control only. Do not use for any other purpose.

NOTE: The so-called "rubber" and "glue" in dwellings, etc. Do not allow children or cooking utensils to be in contact with treated surfaces.

To Control Household Pests, use base oil to meet necessary dilution.

Spray Dilution

Gal
fini
spr.

*2 tablespoons
Cockroaches (1)
spray or apply
behind and be-
hind trails and point
For heavy cock-
roach infestation
monthly applica-
tion. Carpet Beetles
carpeting, and
insects are seen
Spiders and Sc
Make spot appli-
cation. Pantry Pests—e
Drugstore Beet



NON[®] OS-4 INSECTICIDE

FOR PEST CONTROL OPERATORS ONLY.
USE WITH INSECTICIDE BASE OILS.
Keep Out of Reach of Children
SEE LABEL FOR ADDITIONAL CAUTIONS.

..... (propyl-6-methyl-4-pyrimidinyl)	48.7%
..... derivative solvent	44.0%
.....	7.3%
Total:	100.0%

NON OS-4 contains 4 lbs. Diazinon per gallon.
Registered trademark of CIBA-GEIGY CORP.
U.S. Patent No. 2,754,243

Do not make any representation, expressed or implied, concerning the use of this product other than that stated on the label. Beware of the risk of use and/or handling of this material when such instructions are not followed.

CG-8-74

EPA EST No. 148-1L-1

**THOMPSON HAYWARD
CHEMICAL COMPANY**

3333 KANSAS CITY KANSAS 66110

GENERAL INFORMATION

DIAZINON OS-4 is an oil solution containing 4 lbs. of Diazinon per gallon. It is intended for use by pest control operators after dilution with insecticidal base oils for the control of those insects listed below. DIAZINON OS-4 is not intended for use in the manufacture of finished products for resale.

NOTE: The solvent used in this formulation is a petroleum distillate which may stain certain plastic, rubber, and asphalt materials such as tiles and floor coverings. Do not treat such materials. When used in dwellings, care should be taken to avoid deposits which could be frequently contacted by children. Do not allow children in treated areas until surfaces are dry. Do not contaminate food, food containers, or cooking utensils.

DIRECTIONS FOR USE

To Control Household Insects: ** Dilute DIAZINON OS-4 with deodorized kerosene or other suitable base oil to make ½% or 1% Diazinon spray and apply as indicated below. Repeat application as necessary.

Spray Dilution Chart

Gals. of finished spray	Amount of DIAZINON OS-4 to use for	
	½% spray	1% spray
1	1.0 fl. oz.*	2.0 fl. oz.
3	3.0 fl. oz.	6.0 fl. oz.
10	10.0 fl. oz.	20.0 fl. oz. (1.25 pts.)
25	25.0 fl. oz. (1.6 pts.)	50.0 fl. oz. (3.1 pts.)

*2 tablespoons = 1 fl. oz.

Cockroaches (Waterbugs), Ants and Silverfish: Make spot applications of ½% Diazinon as a coarse spray or apply with a paint brush to baseboards, areas around water pipes, cracks and crevices, surfaces behind and beneath cabinets, refrigerators, sinks, stoves and similar areas where these insects hide. Ant trails and points of entry to the premises should also be treated.

For heavy cockroach infestations, make the first application at a strength of 1% Diazinon, followed by monthly applications of ½% Diazinon.

Carpet Beetles: Apply a ½% Diazinon spray in spot applications along baseboards and edges of carpeting, under carpeting and rugs and furniture, in closets and on shelving, and wherever these insects are seen or suspected.

Spiders and Scorpions: Apply a ½% Diazinon spray around windows, doors, and along baseboards. Make spot applications to other areas over which these pests may crawl.

Pantry Pests—exposed stages. (Saw-toothed Grain Beetle, Flour Beetle, Rice Weevil, Cigarette Beetle, Drugstore Beetle, Indian meal Moth)