

NET WEIGHT

POUNDS



Avoid inhalation of dust. Avoid contact with skin.

Avoid contamination of feed and foodstuffs. Some fruits and other plants are susceptible to injury from sulfur under certain climatic conditions. The user is advised not to use sulfur on any crop unless local use has proven that sulfur is safe in that locality.

STORE IN A COOL, DRY PLACE — PROTECT FROM EXCESSIVE HEAT
Burn bag immediately when empty. Stay out of smoke. Do not apply to or allow to drift onto sulfur-sensitive crops or varieties. Do not apply during periods of extreme heat.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label."

DIRECTIONS FOR USE

CAPTAN SULFUR 10-50 DUST is a fungicidal dust formulation adapted for use by ground duster or airplane.

CELERY (Field & Seed Bed only): Pink Rot, Damp-Off. Apply 15 to 20 lbs. per acre. Apply when all seeds have germinated. Repeat at 7 to 10 day intervals as long as disease is active.

GRAPES: Botrytis Rot (Bunch Rot), Powdery Mildew, Silver Mite, Storage Rots. Apply 40 to 50 lbs. per acre. Make 2 applications before bloom and 1 application just after bloom. Repeat periodically, making up to 3 cover applications before bunches close. See re-entry cautions below.

PEACHES: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite, Peach Scab. Apply 50 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest applications when necessary. See re-entry cautions below.

PLUMS: Brown Rot, Storage Rots, Powdery Mildew, Silver Mite. Apply 45 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application when necessary.

RE-ENTRY GRAPES, PEACHES: Do not allow workers to enter treated areas within 1 day (24 hrs.) of application to engage in any activity requiring substantial contact with treated foliage.

POTATOES: Early and Late Blight. Apply 25 to 50 lbs. per acre. Begin application when plants are 2 to 4 inches high and repeat at 7 to 10 day intervals as long as disease threatens.

STRAWBERRIES: Botrytis Rot, Powdery Mildew. Apply 30 lbs. per acre immediately after picking. Repeat application after picking as necessary.

TOMATOES: (Established Plants): Early and Late Blight, Anthracnose, Russet Mite. Apply 25 to 30 lbs. per acre on young plants, increasing to 30 to 50 lbs. per acre on mature plants. Apply at first sign of insects and diseases. Repeat at 5 day intervals as necessary for control. Because of sulfur content of this dust, first obtain permission from your canner or processor before using it on tomatoes intended for sale to such canner or processor.

ACCEPTED
JUL 19 1974
UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 148-1153 SUBJECT
TO APPLICABLE COMMENTS.

CAPTAN* SULFUR 10-50 DUST

ACTIVE INGREDIENTS:

Captan* 10.0%
Sulfur 50.0%

INERT INGREDIENTS:

..... 40.0%
Total 100.0%

*N-trichloromethylthio-4-cyclohexene-1, 2 dicarboximide

CAUTION Keep Out of Reach of Children

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risks of use and/or handling of this material when such use and/or handling is contrary to label instructions.

EPA Reg. No. 148-1153

CG-6-74A



THOMPSON HAYWARD
CHEMICAL COMPANY

P.O. BOX 2383 KANSAS CITY, KANSAS 64116