



### DIRECTIONS FOR USE

This product is formulated for Application by ground or air equipment

**ALMONDS:** Brown Rot, Shot Hole, Scab, Leaf Blight—Use 25-50 lbs per acre. Apply at popcorn, bloom, petal fall stages and up to 5 weeks after petal fall. May be used up to 12 days of harvest if hulls are not fed to dairy animals or animals being finished for slaughter.

**APPLES:** Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Rot, Botryosphaeria, Black Pox—20-35 lbs per acre. Make first application at delayed dormant period. Repeat applications for each infection period or at weekly intervals as long as necessary.

**APRICOTS:** Brown Rot, Jacket Rot—50 lbs per acre. Apply at red bud stage and repeat at 75 percent petal fall.

**BEANS AND PEAS:** Damp-off on beans, root rot on beans and blight on peas—50 to 60 lbs per acre. Work into top 3 or 4 inches of soil before planting.

**CANTALOUPE, CUCUMBERS, SUMMER SQUASH, WINTER SQUASH, WATERMELON, PUMPKINS:** Downy Mildew, Angular Leaf Spot, Anthracnose—Apply 20 to 25 lbs per acre on young plants, increasing to 30 lbs per acre on mature plants. Apply at first sign of disease and continue at weekly intervals as necessary.

**CARROTS:** Septoria Leafspot, Early Blight (Cercospora Blight)—25 lbs per acre. Apply at first sign of disease. Repeat at 6 to 10 day intervals.

**CELERY:** Pink Rot, Damp-off Diseases (field)—20 lbs per acre (seed bed)—40 to 45 lbs per acre. Apply when all seeds have germinated. Repeat as necessary.

**CHERRIES:** Cherry Leaf Spot, Brown Rot—40 to 50 lbs per acre. Apply at popcorn, blossom, and petal fall stages, continue at 10 to 14 day intervals.

**GRAPES:** Mold Control on fruit on drying trays in field—*Penicillium* spp., *Aspergillus* spp., *Botrytis Cinerea*, *Rhizopus* spp.—10 lbs per acre. Apply to fruit on trays for protection of fruit and prevention of above molds. Apply to give even coverage. During damp weather make a second application after trays are turned. Apply by hand, ground duster or airplane. If applied by plane increase to 15 lbs per acre. Do not apply within 5 days of a pre-harvest application of Captan.

**ORNAMENTALS:** Roses—Black Spot, Carnations—Rust, Leaf Spot, Chrysanthemums—Flower Blight—Apply at first sign of disease to give even coverage of upper and lower leaf surfaces. Repeat at 7 to 10 day intervals as necessary.

**PEACH DISEASES:** Brown Rot, Peach Scab—40 to 50 lbs per acre. Make 2 to 3 prebloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make preharvest application where necessary.

**PEARS:** Pear Scab (except Pacific Northwest)—20 to 30 lbs per acre. Make first application during early finger and petal fall stages. Repeat applications for each infection period or at weekly intervals as long as necessary. Russeting may be reduced on Bosc Pears. Do not use on D'Anjou Pears.

NET WEIGHT

POUNDS

# CAPTAN\* 10 DUST

### ACTIVE INGREDIENT:

Captan*	10.0%
INERT INGREDIENTS:	90.0%
Total	100.0%

\*N-trichloromethylthio-4-cyclohexene-1,2 dicarboximide

## CAUTION Keep Out of Reach of Children

See other precautions on side panel.

*May 25, 1974*  
*198-11-27*



THOMPSON MAYNARD  
CHEMICAL COMPANY

**POTATOES:** (seed pieces) Fusarium Seed Rot, Verticillium wilt and damp-off—Dust at 8 ounces per bushel cut seed pieces. (12 ozs. to 100 lbs. cut seed pieces) Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting.

If planting is to be delayed more than 1 or 2 days, the treated pieces should be stored for 2 to 3 days in open crates before bagging.

**POTATOES:** Early and Late Blight—Young Plants—20 to 30 lbs. per acre. Mature Plants—40 lbs. per acre. Begin applications when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

**RASPBERRIES:** Fruit Rot—20 lbs. per acre making applications 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days after mid-harvest.

**SOIL AND GREENHOUSE BENCH TREATMENT:** Preplanting for damp-off and Root Rot Diseases on seedling or transplants of Roses (and other shrubs and trees), many flowers and vegetables (such as beans, peas, cucurbits) and lawn seed beds.—2 lbs. p 1000 sq. ft. (50 to 60 lbs. per acre). Work into upper 3 or 4 inches of soil before planting.

**STRAWBERRIES:** Botrytis Rot—40 lbs. per acre. Make first application when new growth starts in Spring before fruit starts to form. Repeat applications weekly. Severe infection conditions—continue through harvest period, treating immediately after each picking.

**TOMATOES:** Early Blight, Late Blight, Anthracnose, Septoria Leaf Spot—Young Plants—20 to 30 lbs. per acre. Mature Plants—Increase to 30 to 40 lbs. per acre. Apply at first sign of diseases. Repeat at 7 to 10 day intervals.

### CAUTION

Avoid inhalation of Dust. Avoid contact with skin. Do not store or transport with feed or food. Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

This product is toxic to fish. Keep out of lakes, streams, or ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on label.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH CAUTIONS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER. BURN BAG IMMEDIATELY WHEN EMPTY. STAY OUT OF SMOKE. STORE IN COOL, DRY PLACE—PROTECT FROM EXCESSIVE HEAT.

### NOTICE

Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.