

# CAUTION

May be absorbed through skin. Do not breathe spray mist. Do not get on skin, eyes or clothing. Wash thoroughly after handling. Harmful if swallowed.

Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans, pets or livestock.

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams, or ponds. Do not apply where run-off is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

**CAUTION:** Use only clean equipment with adequate mechanical agitation. Do not combine with sulfur or sulfur containing products. Do not use 60 days before or until 90 days after an application of sulfur. Do not apply when temperatures are below 40° or above 90° or when nighttime temperatures fall below 32°. Do not apply when trees are weakened or under moisture stress.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

# SUPERIOR OIL-DIAZINON®

## ACTIVE INGREDIENTS:

Paraffinic hydrocarbon oil<sup>2</sup> 93.00%  
0,0-Diethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 4.50%

## INERT INGREDIENTS:

2.50%  
100.00%

Minimum unsulfonated residue 92% — Viscosity  
90 SSU @ 100° F.

Diazinon is a registered trademark of Ciba-Geigy Corporation.

CONTAINS 1/3 lb. Diazinon/Gallon

**CAUTION Keep Out of Reach of Children**

SEE OTHER PANEL FOR ADDITIONAL CAUTIONS.

crushing and burying in a safe place away from water supplies.

## DIRECTIONS FOR USE

CROP	PEST	DOSAGE PER 100 GALLONS
<b>APRICOTS ALMONDS CHERRIES NECTARINES PEACHES PLUMS PRUNES</b>	Brown Almond Mite Eggs European Red Mite Eggs San Jose Scale Parlatoria Scale Black Cherry Aphid Eggs Forbes Scale Peach Twig Borer Crawler State of Brown. Apricot, Black, San Jose Parlatoria & Oleander Scales.	For Dormant or Delayed Dormant Use 1 1/2 to 2 Gallons

Do not allow workers to reenter treated peaches or nectarines to engage in any activity requiring substantial contact with treated foliage for 5 days.

<b>WALNUTS</b>	San Jose Scale	For Dormant use 5 Gallons per acre in 200 gallons of water
Do not apply after husks open.		

<b>APPLES PEARS</b>	Rosy Apple Aphid Eggs Clover Mite Eggs European Red Mite Eggs San Jose Scale Lecanium Scale Brown Apple Scale Pear Psylla Blister Mite Scale Crawler Tentative Leaf Miner Apple Pear Mite Eggs	For Dormant or Delayed use 1 1/2 to 2 gallons.
-------------------------	---	--

## NOTICE

**Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use.**