



DIAZINON 4 DUST

ACTIVE INGREDIENT:

O,O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate

4%

INERT INGREDIENTS:

96%

Total

100%

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after use. Avoid contact with food, feed, and water. Keep out of reach of children and pets.

Fish and Wildlife Caution

Do not apply to water. Do not apply to areas where fish and wildlife are present. Do not apply to areas where birds are present. Do not apply to areas where bees are present. Do not apply to areas where honey bees are present. Do not apply to areas where bees are present.

Do not apply to areas where fish and wildlife are present. Do not apply to areas where birds are present. Do not apply to areas where bees are present. Do not apply to areas where honey bees are present. Do not apply to areas where bees are present.

Do not apply to areas where fish and wildlife are present. Do not apply to areas where birds are present. Do not apply to areas where bees are present. Do not apply to areas where honey bees are present. Do not apply to areas where bees are present.

Do not apply to areas where fish and wildlife are present. Do not apply to areas where birds are present. Do not apply to areas where bees are present. Do not apply to areas where honey bees are present. Do not apply to areas where bees are present.



P.O. BOX 1200, KANSAS CITY, KANSAS 65110

WORKER RE-ENTRY: Do not allow workers to enter field to engage in any activity requiring substantial contact with treated foliage for 5 days after application.

DIRECTIONS FOR USE

BEANS: Bean Aphid, Leaf Miner, Cucumber Beetle, Leafhopper. Apply 30-35 lbs. per acre. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER: Imported Cabbage Worm, Aphid, Diamond-back moth. Apply 30-35 lbs. per acre. Do not apply within 7 days of harvest for Cabbage. Do not apply within 5 days of harvest for Broccoli and Cauliflower.

CORN: Corn Earworm, Grasshopper. Apply 30-40 lbs. per acre. Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days after treatment.

GRAPES: Leafhopper, Grape Berry Moth, Grape Leaf Folder. Apply 20-25 lbs. per acre. Do not apply within 30 days of harvest. See Re-entry cautions above.

STRAWBERRIES: Aphid. Apply 20-30 lbs. per acre. Do not apply within 7 days of harvest.

TOMATOES: Aphid, Leaf Miner. Apply 30-40 lbs. per acre. Do not apply within 7 days of harvest.

FIGS: Fig Canker, Fig Leaf Fly. Apply 40 lbs. per acre. Do not apply later than 30 days before harvest.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.