



DIAZINON 4 DUST

Under the Fungicide, or as amended, registered under EPA Reg. No.

ACTIVE INGREDIENT:		
O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate	4%
INERT INGREDIENTS:	96%
Total	100%

CAUTION: Keep Out of Reach of Children

Harmful if swallowed. May be absorbed through skin. Do not breathe dust. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of food and feed. Keep out of reach of domestic animals. Do not use on humans or household pets.

Fish and Wildlife Caution

This product is toxic to fish, and wildlife. Birds feeding on treated areas may be killed. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not reuse container. Destroy it by burying or burning. Avoid smoke or fumes.

Diazinon® trademark of CIBA-GEIGY

U. S. Patent No. 2,754,243

EPA Reg. No. 148-1212

CG-7-75

EPA Est. No. 148-CA-1



THOMPSON HAYWARD
CHEMICAL COMPANY

P O BOX 2383 KANSAS CITY KANSAS 66110

WORKER RE-ENTRY: Do not allow workers to enter field to engage in any activity requiring substantial contact with treated foliage for 5 days after application.

DIRECTIONS FOR USE

BEANS: Bean Aphid, Leaf Miner, Cucumber Beetle, Leafhopper. Apply 30-35 lbs. per acre. Do not apply within 7 days of harvest.

BROCCOLI, CABBAGE, CAULIFLOWER: Imported Cabbage Worm, Aphid, Diamond-back moth. Apply 30-35 lbs. per acre. Do not apply within 7 days of harvest for Cabbage. Do not apply within 5 days of harvest for Broccoli and Cauliflower.

CORN: Corn Earworm, Grasshopper. Apply 30-40 lbs. per acre. Do not apply within 2 days of harvest. Do not feed treated forage to animals being finished for slaughter within 2 days after treatment.

GRAPES: Leafhopper, Grape Berry Moth, Grape Leaf Folder. Apply 20-25 lbs. per acre. Do not apply within 10 days of harvest. See Re-entry cautions above.

STRAWBERRIES: Aphid. Apply 20-30 lbs. per acre. Do not apply within 5 days of harvest.

TOMATOES: Aphid, Leaf Miner. Apply 30-40 lbs. per acre. Do not apply within 3 days of harvest.

FIGS: For control of Fruit Fly. Apply 40 lbs. per acre. Do not apply later than 5 days before harvest.

NOTICE: Seller makes no warranty, expressed or implied, concerning the use of product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.