

# DURSBAN

## 1% granular

Under the Federal Insecticide, Fungicide, and Plant Disease Act of 1947

**ACTIVE INGREDIENT:**  
Chlorpyrifos O,O-d-ethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate ..... 1%

**INERT INGREDIENTS:** ..... 99%

100%

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

\*DURSBAN\* — Trademark of the Dow Chemical Company  
U. S. Patent No. 3,244,586

EPA Reg. No. 14C-1267      CG-7-78      EPA Est. No. 148-

**NET WEIGHT 50 POUNDS (22.68 kilograms)**



**THOMPSON-HAYWARD  
CHEMICAL COMPANY**

**P.O. BOX 2383, KANSAS CITY, KANSAS 66110**

**PRECAUTIONARY STATEMENTS**

**HAZARD TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** MAY BE HARMFUL IF SWALLOWED. Do not take internally. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

**NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

**ENVIRONMENTAL HAZARD**

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply where runoff is likely to occur. Use this product only as specified on this label.

**DIRECTIONS FOR USE**

**GENERAL CLASSIFICATION**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**STORAGE:** Keep away from food, feedstuffs, and domestic water supplies. Keep container closed and out of reach of children and pets.

**DISPOSAL:** Do not reuse empty container. Wrap container and put in trash collection.

**LAWNS AND TURF**

Ants, brown dog ticks, chinch bugs, crickets, cutworms, earwigs, sowbugs, pillbugs, sod webworms - 100 = per acre.  
Hyperodes weevil - 200 = per acre.  
European chafer, Japanese beetle larvae - 200-400 = per acre.

Apply DURSBAN 1% GRANULAR uniformly using any hand or power operated gravity or rotary type spreader designed for application of granular materials. Treat when insects first appear. Water in immediately. Do not mow for 24 hours following treatment.

For smaller than one acre, see conversion chart below.

AREA TO BE TREATED	POUNDS DURSBAN 1% GRANULAR		
	100 = A rate	200 = A rate	400 = A rate
1000 sq. ft.	2	5	10
5000 sq. ft.	12	25	50

This product may be used to control the insects listed outdoors around domestic buildings, medical facilities and schools, to control the infestation and reinfestations of buildings. Apply uniformly at rates listed in a band 6 to 10 feet wide around and adjacent to building. Water into soil. Do not mow for 24 hours following treatment. Do not apply to D. chondria.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.