



ACTIVE INGREDIENTS:	By Weight
Carbaryl (1-naphthyl N-methylcarbamate)	10%
INERT INGREDIENTS	90%
TOTAL	100%

SEVIN is the registered trade mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

If swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching finger to back of throat. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.
 If inhaled: Remove victim to fresh air. Apply respiration if indicated.
 If on skin: Wash immediately with soap and water.
 If in eyes: Flush eyes for at least 15 minutes with water. Call a physician immediately.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: MAY BE HARMFUL IF SWALLOWED. Do not take internally. Avoid contact with eyes, skin and clothing. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention if irritation persists. Keep children and pets off treated areas until this material is washed into the soil and grass is dry.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ENVIRONMENTAL HAZARD

This product is toxic to fish, birds and other wildlife. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label. Shrimp and crab may be killed at application rates recommended on this label. Do not apply this product where these are considered important resources.

Bee Caution: May kill honeybees in substantial numbers. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not use when value of bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

Avoid contamination of Food, Feed, Feeding troughs and Watering Receptacles.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide Disposal: Pesticides, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable Federal, state or local procedures.

Container Disposal: Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment. Then dispose of bags in a sanitary landfill or by incineration if allowed by State and local authorities.

EPA Reg No. 148- EPA Est. No. 9859-EL-1
NET WEIGHT 50 POUNDS (22.68 kilograms)



P.O. BOX 2383, KANSAS CITY, KANSAS 66110

Frank G. ...

12/10/60

1502
148-1276

17

PEST CONTROL

DOMESTIC DWELLINGS -- Earwigs - 5 to 7½ lbs. per 5000 square feet. Apply to soil surface, mulch, and debris around moist shady places. Fleas - 2½ to 5 lbs. per 5000 square feet. Apply liberally to pens, yards, lawns, and under buildings around the home. Force the dust into cracks and crevices. Repeat as needed. Millipedes - 5 to 7½ lbs. per 5000 square feet. Apply to soil surface, mulch, and debris around moist shady places. Mosquitoes - 2½ to 5 lbs. per 5000 square feet. Apply to lawns, shrubbery, and other vegetation around the home. Repeat as needed. Ticks -- 1½ to 2½ lbs. per 5000 square feet. Apply liberally to ground, debris and vegetation to a height of about 2 feet on lawns, vacant fields, and wooded areas around the home. Repeat as needed.

ALL RATES SHOULD BE APPLIED AS FOLIAGE APPLICATIONS UNLESS OTHERWISE NOTED.

ORNAMENTALS

LAWNS AND TURFS -- Ants, Chinch bugs, Crickets, Cutworms, Earwigs, Fall army-worm, Fleas, Leafhoppers, Millipedes, Mosquitoes, Sod webworms, Ticks -- 10 lbs per 5000 square feet. For best results, mow lawn, remove clippings and apply immediately after rain or watering. Water following treatment. Keep children and pets off lawn until material is watered in and grass is dry.

HERBACEOUS PLANTS AND BULBS (Annual plants-Herbaceous plants). Sevin injures Boston ivy and Virginia creeper. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity. Blister beetles, Flea beetles, Japanese beetle, Lacebugs, Leafhoppers, leafrollers, Plant bugs, Psyllids, Rose aphid, Thrips (exposed) -- 15 to 20 lbs. per acre.

WOODY SHRUBS, TREES AND VINES -- Sevin injures Boston ivy and Virginia creeper. Injury to tender foliage may occur if plants are wet when treated or in presence of high humidity. Apple aphid, Bagworm, Birch leafminer, Boxelder bug, Boxwood leafminer, Elm leaf aphid, Elm leaf beetle, Gypsy moth, Japanese beetle, Lace bugs, Leafhoppers, Leafrollers, Oak leafminers, Orange tortrix, Periodical cicada, Rose aphid, Roseslug, Scales, Thrips (exposed), Tent caterpillars, Willow leaf beetles -- 15 to 20 lbs. per acre. Plant bugs, Psyllids -- 15 to 20 lbs. per acre.

NONAGRICULTURAL LANDS, PASTURE, RANGELAND -- Mosquitoes -- 2½ to 5 lbs. per acre. Air blast duster. Thorough coverage is necessary for effective control. For optimum control, apply in early morning or evening when air is calm and adults are active. May kill shrimp and crabs. Do not use in areas where these are important resources. Mosquitoes -- 2½ to 10 lbs. per acre with low pressure hydraulic ground duster. For optimum control, apply in early morning or evening when air is calm and adults are active. May kill shrimp and crabs. Do not use in areas where these are important resources.

WOODED AREAS -- House flies, Mosquitoes, Sand flies -- 3 lbs. per acre. Apply with power duster or large hand duster.

URBAN AND RURAL OUTDOOR AREAS (PUBLIC HEALTH) Plague reservoir host animals -- For use only by or under the supervision of State or Federal public health or wildlife officials. Fleas -- 1 oz. per burrow. Apply by hand or power applicator. Fleas -- 2 oz. per bait station. Use 4-6 bait stations per acre. Fleas -- 10 lbs. per acre. Apply by hand or power applicator as an area dust treatment. Consult your State Fish and Game Agency before applying this product.

DOG HOUSES, KENNELS, SLEEPING QUARTERS -- 1½ to 2½ lbs. per 5000 square feet for ticks. 2½ to 5 lbs. per 5000 square feet for fleas. Apply liberally by garden-type duster. Force dust into cracks and crevices. Repeat as needed. Do not expose kittens or puppies less than 4 weeks of age to treated areas.

Back - Daniel

